



P: 1300 688 522
E: info@nutripath.com.au
A: PO Box 442 Ashburton VIC 3142

Date of Birth : 10-Nov-1981
Sex : F
Collected : 12/Jul/2021
Received: 16/Jul/2021
35A DAY ROAD
NORTHGATE QLD 4013
Lab id : **375411** UR#: 6587824

6 EDWARDS BAY ROAD
MOSMAN NSW 2088

BIOCHEMISTRY

BLOOD - SERUM

25 OH VITAMIN D

Result	Range	Units
61	50 - 250	nmol/L



Vitamin D Comment

The Working Group of the Australian and New Zealand Bone and Mineral Society, Endocrine Society of Australia and Osteoporosis Australia have recommended that the Vitamin D level should be greater than 50 nmol/L.

Recommended Target Range:	125 - 200	nmol/L
Deficient Level:	< 50	nmol/L
Insufficient Level:	50 - 74	nmol/L
Sufficient Level:	75 - 250	nmol/L
Excessive Level:	> 250	nmol/L

COMMENT:

25-hydroxy Vitamin represents the body's stores/reserves level. Where levels are deemed sub-optimal, consider assessing 1-25-dihydroxy Vitamin D3 (the active form of the molecule). This may be tested off the existing specimen at the laboratory within 7 days. If you wish to proceed with this test, please forward a new request form for this patient.

BLOOD - PLASMA

HOMOCYSTEINE

Result	Range	Units
4.0 *L	5.0 - 12.0	umol/L



INTEGRATIVE MEDICINE

BLOOD - SERUM

Free Copper Index

COPPER

Result	Range	Units
19.0	11.0 - 22.0	umol/L



CAERULOPLASMIN

Result	Range	Units
0.21	0.18 - 0.40	g/L



% Free Copper

Result	Range	Units
48 *H	5 - 25	%



(*) Result outside normal reference range

(H) Result is above upper limit of reference range (L) Result is below lower limit of reference range





P: 1300 688 522
 E: info@nutripath.com.au
 A: PO Box 442 Ashburton VIC 3142

Date of Birth : 10-Nov-1981
 Sex : F
 Collected : 12/Jul/2021
 Received: 16/Jul/2021
 35A DAY ROAD
 NORTHGATE QLD 4013
 Lab id : **375411** UR#: 6587824

6 EDWARDS BAY ROAD
 MOSMAN NSW 2088

Integrative Medicine Comments

Plasma Copper Reference Ranges (Additional):

Female on Contraceptives: 20.8 - 34.8 umol/L

ELEVATED Cu/Zn RATIO:

The ratio of copper to zinc is clinically more important than the individual levels of each analyte.

Elevated copper/Zinc ratios can be particularly serious for persons with low blood histamine (over methylation). This combination of imbalances has been associated with anxiety, panic disorders, paranoia and hallucinations.

ELEVATED FREE COPPER:

Consider chelation therapy, Vit C, Zinc, Molybdenum, Glutamine, Histidine and threonine.

Optimal free Copper of 5-25% is recommended by Dr B Walsh.

Retesting Zinc and Copper levels after 3 months of treatment is recommended.

Signs of Cu overload include hyperactivity, skin sensitivity to metals. Skin tags, estrogen dominance, emotional meltdowns, tinnitus, abnormal periods.

HISTAMINE

19.0 *L 28.0 - 51.0 ug/L



(*) Result outside normal reference range

(H) Result is above upper limit of reference range (L) Result is below lower limit of reference range



P: 1300 688 522
E: info@nutripath.com.au
A: PO Box 442 Ashburton VIC 3142

Date of Birth : 10-Nov-1981
Sex : F
Collected : 12/Jul/2021
Received: 16/Jul/2021
35A DAY ROAD
NORTHGATE QLD 4013
Lab id : **375411** UR#: 6587824

6 EDWARDS BAY ROAD
MOSMAN NSW 2088

Histamine Comment

LOW WB HISTAMINE LEVELS:

Low Whole Blood Histamine is otherwise known as Histapenia.

In assessing Histamine levels, Diamine Oxidase (DAO) should also be tested concurrently.

Serum/plasma copper levels in histapenic patients are often abnormally high. As copper is a brain stimulant (and also destroys histamine), the elevated serum/plasma (and presumably brain) copper level probably accounts for many visible symptoms, including the low blood histamine level.

Behavioral symptoms in high-copper histapenia include paranoia and hallucinations in younger patients. In older patients, depression may predominate. Some studies of schizophrenics have revealed high blood copper, as seen in histadelia, with low urinary copper (showing that copper is being retained) as well as low blood zinc.

Treatment considerations:

Treatment protocol consists of the administration of Amino Acid Histidine, (converts to histamine), a diet high in protein, supplement with B6, Folic Acid & B12 injections as they increase histamine levels while lowering the degree of symptoms. Zinc and manganese with vitamin C remove copper from the tissues. Copper destroys histamine and therefore as copper levels decrease, histamine levels should return towards normal. With this treatment the high blood copper is slowly reduced and symptoms are slowly relieved in several months' time.

Treating histadelia is not a quick process. Early noticeable results/improvements may not be seen until close to 6-10 weeks. Treatment can take more than 12 months to complete.

To find out more about this see Mental Illness: The Nutrition Connection, a book by Carl Pfeiffer. The prognosis is seen to be good if the histadelic patient cooperates with treatment and works to give up detrimental addictions.

ZINC	13.7	9.0 - 19.0	umol/L	
Copper/Zinc Ratio	1.39 *H	0.80 - 1.00	RATIO	

Tests ordered: CFee,IMPEI,FCuInd,VITD,ZN,Cu/Zn,HIAM,HOMO

(*) Result outside normal reference range

(H) Result is above upper limit of reference rang (L) Result is below lower limit of reference range