



P: 1300 688 522
E: info@nutripath.com.au
A: PO Box 442 Ashburton VIC 3142

Date of Birth : 10-Apr-1981
Sex : F
Collected : 6/Apr/2022
Received: 12/Apr/2022
14 WOODVALE AVENUE
EPPING NSW 2121
Lab id : **3809021** UR#: 6603470

6 EDWARDS BAY ROAD
MOSMAN NSW 2088

Vaginal Microbiome Profile

Vaginal pH.

4.5

3.5 - 4.5



Opportunistic Bacteria	Result	Range	Units	
Enterococcus faecalis:	<DL	< 1.0	x10 ⁵ CFU/ml	
Escherichia coli:	<DL	< 1.00	x10 ⁵ CFU/g	
Klebsiella pneumoniae:	<DL	< 1.00	x10 ⁵ CFU/ml	
Proteus mirabilis:	<DL	< 1.00	x10 ⁵ CFU/ml	
Pseudomonas aeruginosa:	<DL	< 1.00	x10 ⁵ CFU/ml	
Streptococcus agalactiae:	<DL	< 1.00	x10 ⁵ CFU/ml	
Staphylococcus aureus:	<DL	< 1.00	x10 ⁵ CFU/ml	
Gardnerella vaginalis:	9.07 *H	< 1.00	x10 ⁵ CFU/ml	
Atopobium vaginae:	7.56 *H	< 1.00	x10 ⁵ CFU/ml	
Prevotella species:	<DL	< 1.00	x10 ⁵ CFU/ml	
Megasphaera species:	<DL	< 1.00	x10 ⁵ CFU/ml	
Ureaplasma species	0.13	< 1.00	x10 ⁶ CFU/ml	
Mycoplasma species	<DL	< 1.00	x10 ⁶ CFU/ml	

Sexually Transmitted Infections

Trichomonas vaginalis:	Not Detected
Chlamydia trachomatis:	Not Detected
Neisseria gonorrhoeae:	Not Detected
Herpes Simplex Virus-1:	Not Detected
Herpes Simplex Virus-2:	Not Detected

COMMENT:

Not Detected results indicate the absence of detectable DNA in this sample. A negative result does not completely exclude infection.

Opportunistic Fungal pathogens

Candida albicans:	<DL	< 1.00	x10 ⁵ CFU/ml	
Candida glabrata:	<DL	< 1.00	x10 ⁵ CFU/ml	
Candida krusei:	<DL	< 1.00	x10 ⁵ CFU/ml	
Candida parapsilosis:	<DL	< 1.00	x10 ⁵ CFU/ml	
Candida tropicalis:	<DL	< 1.00	x10 ⁵ CFU/ml	

Beneficial Bacteria:

Total Lactobacillus:	0.54 *L	> 1.00	x10 ⁶ CFU/ml	
Lactobacillus crispatus:	0.36 *L	> 1.00	x10 ⁶ CFU/ml	
Lactobacillus gasseri:	<DL *L	> 1.00	x10 ⁶ CFU/ml	
Lactobacillus iners:	<DL *L	> 1.00	x10 ⁶ CFU/ml	
Lactobacillus jensenii:	0.18 *L	> 1.00	x10 ⁶ CFU/ml	
Lactobacillus rhamnosus:	<DL *L	> 1.00	x10 ⁶ CFU/ml	
Lactobacillus salivarius:	<DL *L	> 1.00	x10 ⁶ CFU/ml	
Lactobacillus vaginalis:	<DL *L	> 1.00	x10 ⁶ CFU/ml	

Bacterial Vaginosis:

Bacterial vaginosis Negative





P: 1300 688 522
 E: info@nutripath.com.au
 A: PO Box 442 Ashburton VIC 3142

Date of Birth : 10-Apr-1981
 Sex : F
 Collected : 6/Apr/2022
 Received: 12/Apr/2022
 14 WOODVALE AVENUE
 EPPING NSW 2121
 Lab id : **3809021** UR#: 6603470

6 EDWARDS BAY ROAD
 MOSMAN NSW 2088

Vaginal Microbiome Comments

VAGINAL pH NORMAL:

The typical vaginal pH is 3.5-4.5. Prepubertal and postmenopausal pH levels are normally >5 pH. With the increase of the oestrogen levels around puberty, the genital mucosa thickens and becomes colonized with *Lactobacillus* species which produce lactic acid and hydrogen peroxide to lower the pH below 4.5.

GARDNERELLA VAGINALIS ELEVATED:

Gardnerella is a part of normal vaginal anaerobic flora but overgrowth can cause Bacterial vaginosis. This is a poly-microbial infection which suppresses dominance of normal vaginal *Lactobacillus* spp. (Total *Lactobacillus* <10⁶ CFU/ml), presence of clue cells, alkaline vaginal pH (>4.5) and fishy vaginal discharge. BV may be asymptomatic or cause symptoms such as itching and malodorous discharge (often thin and white/grey). It is associated with an increased risk of preterm delivery, pelvic inflammatory disease and an increased risk of acquisition of sexually transmitted infections.

Can be treated after ruling out allergy/pregnancy status:

Metronidazole 400 mg orally, 12-hourly for 7 days or Metronidazole 0.75% vaginal gel 1 applicatorful intravaginally at bedtime for 5 nights

OR
 Clindamycin 2% vaginal cream 1 applicatorful intravaginally, at bedtime for 7 nights (If pregnant or allergic to metronidazole)

General advice for along with above treatment as follows:

- o Regular salt or warm water only washes (no douching)
- o Good Personal Hygiene
- o Avoid irritants (soaps/perfumes)
- o Use barrier protection during sex

ATOPOBIUM VAGINAE ELEVATED:

Atopobium vaginae is a gram-positive facultative anaerobic bacterial species that may be involved in BV in some women as a result of a disruption in the normal vaginal microflora.

BV is typically a polymicrobial infection characterized by disruption in *Lactobacillus* dominance (Total *Lactobacillus* <10⁶ CFU/ml), increased pH (>4.5) and the presence of mainly anaerobic microorganisms including *Atopobium vaginae* (>10⁵ CFU/ml).

LACTOBACILLUS:

Lactobacillus is the predominant genus in a healthy vaginal microbiota, and functions to inhibit the adhesion and proliferation of opportunistic and primary pathogens.

The presence of different *Lactobacillus* species is a major factor in the stability of the vaginal microbiome. Women with *L. iners*-dominant microbiomes are more likely to harbor *Candida* than women with *L. crispatus*-dominant microbiomes (due to higher production of lactic acid by *L. crispatus* compared to *L. iners*), leading to better anti-*Candida* activity (impeding *Candida* colonization) than *L. iners* through a greater production of lactic acid. Furthermore, *L. iners* dominance has been associated with other negative health outcomes such as increased risks of Chlamydia trachomatis infection, incident Bacterial Vaginosis and defects in vaginal mucus that compromise antiviral barrier function.

TOTAL LACTOBACILLUS LEVELS LOW:

Total *Lactobacillus* quantification should be >1x10⁶ CFU/ml in a healthy Vaginal Microbiome. Production of H₂O₂ by *Lactobacillus* species is essential in inhibiting the overgrowth of pathogens. In cases where total *Lactobacillus* levels are low, presence of pathogenic bacteria should be reviewed and probiotic therapy should be considered.

Microorganisms not belonging to the *Lactobacillus* genus with the population equal to or greater than 1x 10⁵ CFU/ml is considered to be disturbing the vaginal ecosystem equilibrium.



P: 1300 688 522
E: info@nutripath.com.au
A: PO Box 442 Ashburton VIC 3142

Date of Birth : 10-Apr-1981
Sex : F
Collected : 6/Apr/2022
Received: 12/Apr/2022
14 WOODVALE AVENUE
EPPING NSW 2121
Lab id : **3809021** UR#: 6603470

6 EDWARDS BAY ROAD
MOSMAN NSW 2088

References:

Pacha-Herrera et. al., 2020, Frontiers in Cellular and Infection Microbiology, 10:303.
Oerlemans et. al., 2020, Europe PMC, 10(11).
Tomusiak et. al., 2013, Polish Society of Gynaecologists, 84:352-358.

Sex. Transmitted Infection Comments

TRICHOMONAS VAGINALIS – Not Detected:

This does not completely exclude the possibility of infection as is dependent on an adequate specimen collection. If you have symptoms, please consult with your healthcare practitioner.

CHLAMYDIA TRACHOMATIS – Not Detected:

This does not completely exclude the possibility of infection as is dependent on an adequate specimen collection. If you have symptoms, please consult with your healthcare practitioner.

NEISSERIA GONORRHOEAE – Not Detected:

This does not completely exclude the possibility of infection as is dependent on an adequate specimen collection. If you have symptoms, please consult with your healthcare practitioner.

HERPES SIMPLEX VIRUS Type 1 – Not Detected:

This does not completely exclude the possibility of infection as is dependent on an adequate specimen collection. If you have symptoms, please consult with your healthcare practitioner.

HERPES SIMPLEX VIRUS Type 2 – Not Detected:

This does not completely exclude the possibility of infection as is dependent on an adequate specimen collection. If you have symptoms, please consult with your healthcare practitioner.