

Gavan, Jane  
5 White Road, QUINNS ROCKS. 6030  
**Phone:** 0422973396  
**Birthdate:** 27/07/1951 **Sex:** F **Medicare Number:** 6000 09304 8-2  
**Your Reference:** 2022QU0002907 **Lab Reference:** 2022QU0002907-1  
**Laboratory:** boorarad  
**Addressee:** Dr Anne Chester **Referred by:** Dr Anne Chester

**Name of Test:** XR LEFT FOOT  
**Requested:** 06/05/2022 **Collected:** 06/05/2022 **Reported:** 06/05/2022  
11:00

**X-RAY - LEFT FOOT**

**CLINICAL HISTORY:**

Pain medially.

**FINDINGS:**

Hallux valgus.  
Bones are mildly osteopenic.  
Degenerative osteoarthritis at the TMT and intertarsal joints.  
Hammertoe configuration noted.  
No fracture.  
Small calcaneal spur.  
Lisfranc and subtalar articulations are maintained.

If there are ongoing symptoms, an ultrasound may be considered.

Yours Sincerely,

**Dr Jonathan Waner**  
**MBBCh, FRANZCR**

Copy and paste this link to view images - An HTTP error occurred (12031: Unknown)



Gavan, Jane  
5 White Road, QUINNS ROCKS. 6030  
Phone: 0422973396  
Birthdate: 27/07/1951 Sex: F Medicare Number: 6000 09304 8-2  
Your Reference: 2022QU0003209 Lab Reference: 2022QU0003209-1  
Laboratory: boorabad  
Addressee: Dr Anne Chester Referred by: Dr Anne Chester  
  
Name of Test: US Left Hindfoot  
Requested: 17/05/2022 Collected: 18/05/2022 Reported: 18/05/2022  
12:48

#### ULTRASOUND of LEFT MIDFOOT

**Clinical History:**  
painful left foot

**Comparison:**  
Reference is made to x-ray performed today.

#### Findings:

Targeted scan performed over dorsal and medial midfoot where patient is most symptomatic.

There is global bony irregularity of the intertarsal joints and 2nd to 4th tarsometatarsal joints with synovial thickening and hypervascularity.

Region of interest corresponds to medial talonavicular and naviculo-medial-cuneiform area, demonstrating bony irregularity with surrounding hypervascularity.

The extensor and flexor tendons appear normal. Tibial nerve appears sonographically unremarkable.

#### Conclusion:

Global arthritic synovitis of the intertarsal and tarsometatarsal joints.  
Patient is most symptomatic at the talonavicular and naviculo-medial cuneiform joints.

Thank you for your referral.

Electronically authorised.

Yours sincerely,

Dr Jonathan Waner  
Consultant Radiologist

Sonographer: S.Ooi/ M.Dama

Copy and paste this link to view images - An HTTP error occurred (12031: Unknown)