

P: 1300 688 522 E: info@nutripath.com.au

-.CASEY CLEMENTS 17A CAMERON STREET MACLEAN NSW 2463

TOM HANNA 20-Jan-1984 Male

187 TULLYMORGAN ROAD LAWRENCE NSW 2460

LAB ID : 3853807 UR NO. : 6160935 Collection Date : 17-Nov-2022 Received Date:17-Nov-2022



3853807

MICROBIOLOGY

STOOL, SPOT Result Range Units

Shiga Toxin Summary

SHIGA-LIKE TOXIN E. COLI (STEC): Detected

Method: PCR

Comment:

Shiga-like toxin E.coli is seen in faecal contamination of food (undercooked beef, raw milk, and unpasteurized juice) and water. It may also be seen in farm workers/agricultural families/pet farms. Implicated in haemorrhagic colitis, with fever, abdominal cramping, fatigue, nausea, and frank fresh bloody diarrhoea with no faeces, lasting up to a week. May complicate to haemolytic uremic syndrome (HUS), causing hepato-renal shutdown.

Treatment .

Managed conservatively with supportive therapy, reassurance with warning signs of HUS. B lactam (cell wall active) antibiotics are contraindicated with mild symptoms as can principate HUS, which warrants in-patient IV non B lactam antibiotics.

This test was confirmed by a secondary PCR test.

Dr N. Maheshwari Clinical Pathologist Microbiology

Tests ordered: STEC