



2/5 BIRTWILL STREET, COOLUM BEACH, QLD 4573
Phone: 07 5436 0888 Fax: 07 54433494

To: Dr Physio Synergy Physio
Referred By:
Dr Shauna Vivian
PEREGIAN SPRINGS DOCTORS
RIDGEVIEW DRIVE
PEREGIAN SPRINGS QLD 4573

Ms Denise Donohoe
DOB: 15 January 1957
UR: BAH940Y
Visit #: COL-3434812
Service Date: 28 April 2022

Copies To: Dr [REDACTED]

Visit Description: US BILATERAL HIPS, XRAY LEFT FOOT

BILATERAL HIP/GROIN ULTRASOUND AND XRAY LEFT FOOT

Clinical Indications:

Bilateral hip pain.
Left > Right.
Kicked foot again.? Fracture second toe. Fifth toe fracture in January.

Findings:

ULTRASOUND BILATERAL HIP/GROIN

Right

No joint effusion.

Anterior:

Iliopsoas tendon: Normal.
Iliopsoas bursa: Normal.

Lateral:

Gluteus medius: Complete tear; gap length 18mm.
Gluteus minimus: Complete tear; gap length Full Retraction.
Trochanteric bursa: fluid, focally tender.
Greater trochanter: Normal.
TFL origin: Normal.
ITB: Normal.

Conclusion:

**Gluteus medius and gluteus minimus tendon tears.
Symptomatic trochanteric bursitis.**

Left

No joint effusion.

Anterior:

Iliopsoas tendon: Normal.
Iliopsoas bursa: Normal.

Lateral:

Gluteus medius: Focal tendon thickening/tendinopathy towards the superior posterior facet.
Gluteus minimus: Diffuse tendon thickening/tendinopathy.
Trochanteric bursa: fluid, focally tender.
Greater trochanter: Normal.
TFL origin: Normal.
ITB: Normal

The information contained in this facsimile message is legally privileged and confidential, intended only for the use of the individual named above. If the receiver is not the intended recipient the receiver is hereby notified that any use, dissemination, distribution, publication or copying of this facsimile is prohibited. If you have received this facsimile in error please notify the practice immediately and arrangements will be made to retrieve or destroy it.

Conclusion: Gluteus medius & gluteus minimus tendinopathy
Symptomatic trochanteric bursitis