



3934192

COMPLETE MICROBIOME MAPPING

General Macroscopic Description

	Result	Markers
Stool Colour	Brown	Colour - Brown is the colour of normal stool. Other colours may indicate abnormal gut health.
Stool Form	Formed	Form - Sample form is categorised using the Bristol stool chart. A comment on stool appearance can be found in the comments section.
Mucous	Not Detected	Mucous - Mucous production may indicate the presence of an infection and/or inflammation.
Occult Blood	Negative	Blood (Macro) - The presence of blood in the stool may be the result of several causes besides colorectal bleeding, including hemorrhoids or gastrointestinal infection.


Short Chain Fatty Acids

	Result	Range	Units	
Short Chain Fatty Acids, Beneficial	44.7	> 13.6	umol/g	
Butyrate	19.9	10.8 - 33.5	%	
Acetate	54.7	44.5 - 72.4	%	
Propionate	21.1	0.0 - 32.0	%	
Valerate	4.3	0.5 - 7.0	%	

GIT Functional Markers

	Result	Range	Units	
Calprotectin.	82.0 *H	0.0 - 50.0	ug/g	
Pancreatic Elastase	>800.0	> 200.0	ug/g	
Secretory (slgA)	1318.8	510.0 - 2040.0	ng/mL	
Zonulin	34.5	0.0 - 107.0	ng/mL	
Beta glucuronidase	2825.8	368.0 - 6266.0	U/g	
Steatocrit	<1.0	0.0 - 15.0	%	
a-Transglutaminase IgA	28.0	0.0 - 100.0	units/L	

Microbiome Mapping Summary



Key Phyla Microbiota

Firmicutes:Bacteroidetes Ratio

0.28 < 1.00

RATIO



Relative Commensal Abundance of the 6 Phyla groups can be found on page 4 of this report





3934192

Parasites and Worms.	Result	Range	Units
----------------------	--------	-------	-------

Parasitic Organisms

Cryptosporidium species	<dl	< 1.0	x10 ⁵ org/g	
Entamoeba histolytica.	<dl	< 1.0	x10 ⁵ org/g	
Giardia intestinalis	<dl	< 1.0	x10 ⁵ org/g	
Blastocystis hominis.	<dl	< 1.0	x10 ⁵ org/g	
Dientamoeba fragilis.	<dl	< 1.0	x10 ⁵ org/g	
Endolimax nana	<dl	< 1.0	x10 ⁵ org/g	
Entamoeba coli.	<dl	< 5.0	x10 ⁵ org/g	
Pentatrichomonas hominis	<dl	< 1.0	x10 ⁵ org/g	

Worms

Ancylostoma duodenale, Roundworm	Not Detected			
Ascaris lumbricoides, Roundworm	Not Detected		Necator americanus, Hookworm	Not Detected
Trichuris trichiura, Whipworm	Not Detected		Enterobius vermicularis, Pinworm	Not Detected
Enterocytozoon spp	Not Detected		Hymenolepis spp, Tapeworm	Not Detected
Strongyloides spp, Roundworm	Not Detected		Taenia species, Tapeworm	Not Detected

Comment: Not Detected results indicate the absence of detectable DNA in the sample for the worms reported.

NOTE: Reflex testing is performed on clinically indicated samples

Opportunistic Bacteria/Overgrowth	Result	Range	Units
-----------------------------------	--------	-------	-------

Bacillus species.	<dl	< 1.00	x10 ⁴ CFU/g	
Enterococcus faecalis	<dl	< 1.00	x10 ⁵ CFU/g	
Enterococcus faecium	<dl	< 1.00	x10 ⁵ CFU/g	
Morganella species	<dl	< 1.00	x10 ⁵ CFU/g	
Pseudomonas species	<dl	< 1.00	x10 ⁴ CFU/g	
Pseudomonas aeruginosa.	<dl	< 3.00	x10 ⁴ CFU/g	
Staphylococcus species	<dl	< 1.00	x10 ³ CFU/g	
Staphylococcus aureus	<dl	< 5.00	x10 ³ CFU/g	
Streptococcus species	2.81	< 3.00	x10 ⁶ CFU/g	
Methanobrevibacter smithii	<dl	< 3.50	x10 ⁵ CFU/g	
Desulfovibrio piger	<dl	< 18.00	x10 ⁷ CFU/g	
Enterobacter cloacae complex.	<dl	< 5.00	x10 ⁵ CFU/g	

Potential Autoimmune Triggers

Citrobacter species.	<dl	< 5.00	x10 ⁴ CFU/g	
Citrobacter freundii.	<dl	< 5.00	x10 ⁴ CFU/g	
Klebsiella species	<dl	< 5.00	x10 ³ CFU/g	
Klebsiella pneumoniae.	<dl	< 5.00	x10 ⁵ CFU/g	
Prevotella copri	<dl	< 1.00	x10 ⁹ CFU/g	
Proteus species	<dl	< 5.00	x10 ⁵ CFU/g	
Proteus mirabilis.	<dl	< 1.00	x10 ⁴ CFU/g	
Fusobacterium species	<dl	< 10.00	x10 ⁴ CFU/g	

Fungi & Yeast	Result	Range	Units
---------------	--------	-------	-------

Candida species.	<dl	< 5.00	x10 ⁵ CFU/g	
Candida albicans.	<dl	< 5.00	x10 ⁴ CFU/g	
Geotrichum species.	<dl	< 3.00	x10 ² CFU/g	
Saccharomyces cerevisiae.	<dl	< 3.00	x10 ³ CFU/g	
Rhodotorula species.	<dl	< 1.00	x10 ³ CFU/g	



3934192

Bacterial Pathogens	Result	Range	Units	
Aeromonas hydrophila.	<dl	< 1.00	x10 ³ CFU/g	
Campylobacter species.	<dl	< 1.00	x10 ⁵ CFU/g	
C. difficile, Toxin A	<dl	< 1.00	x10 ⁴ CFU/g	
C. difficile, Toxin B	<dl	< 1.00	x10 ⁴ CFU/g	
Enterohemorrhagic E. coli	<dl	< 1.00	x10 ⁵ CFU/g	
Enteroinvasive E. coli/Shigella	<dl	< 1.00	x10 ³ CFU/g	
Enterotoxigenic E. coli LT/ST	<dl	< 1.00	x10 ⁵ CFU/g	
Shiga-like Toxin E. coli stx1	<dl	< 1.00	x10 ⁴ CFU/g	
Shiga-like Toxin E. coli stx2	<dl	< 1.00	x10 ⁴ CFU/g	
Salmonella species.	<dl	< 1.00	x10 ⁵ CFU/g	
Vibrio species.	<dl	< 1.00	x10 ⁴ CFU/g	
Yersinia species.	<dl	< 1.00	x10 ⁵ CFU/g	
Helicobacter pylori	<dl	< 1.0	x10 ³ CFU/g	

Comment: Helico Pylori virulence factors will be listed below if detected POSITIVE

H.pylori Virulence Factor, babA	Not Detected	H.pylori Virulence Factor, cagA	Not Detected
H.pylori Virulence Factor, dupA	Not Detected	H.pylori Virulence Factor, iceA	Not Detected
H.pylori Virulence Factor, oipA	Not Detected	H.pylori Virulence Factor, vacA	Not Detected
H.pylori Virulence Factor, virB	Not Detected	H.pylori Virulence Factor, virD	Not Detected

Viral Pathogens	Result	Range	Units
Adenovirus 40/41	Not Detected		
Norovirus GI/II	Not Detected		
Rotavirus A	Not Detected		
Sapovirus (I,II,IV,V)	Not Detected		
Astrovirus (hAstro)	Not Detected		

Normal Bacterial GUT Flora	Result	Range	Units	
Bacteroides fragilis	1.1 *L	1.6 - 250.0	x10 ⁵ CFU/g	
Bifidobacterium species	1.0 *L	> 6.7	x10 ⁵ CFU/g	
Bifidobacterium longum	0.7 *L	> 5.2	x10 ⁵ CFU/g	
Enterococcus species	10.6	1.9 - 2000.0	x10 ³ CFU/g	
Escherichia species	2318.0	3.7 - 3800.0	x10 ⁴ CFU/g	
Lactobacillus species	4.0 *L	8.6 - 6200.0	x10 ³ CFU/g	
Lactobacillus rhamnosus	3.2 *L	8.3 - 885.0	x10 ³ CFU/g	
Clostridium species	14.6	5.0 - 50.0	x10 ⁷ CFU/g	
Oxalobacter formigenes	7.63	> 5.00	x10 ⁶ CFU/g	
Akkermansia muciniphila	50.30 *H	1.00 - 50.00	x10 ⁷ CFU/g	
Faecalibacterium prausnitzii	262.0	200.0 - 3500.0	x10 ⁶ CFU/g	

Methodology:

GIT Functional markers performed by GCMS,EIA,FEIA.

Bacteriology,Virology,Fungi,Parasites & Worms performed by PCR,qPCR.







<dl = result below detectable limit. *H = Result greater than the reference range. *L = Result less than the reference range




3934192

Introduction:


Your gut microbiome is a collective name for the 40 trillion cells and up to 1000 microbial species that include bacteria, viruses, fungi, parasites, and archaea and reside in our gut. The number of gut bacterial cells is approximately equal to the total number of human cells in our body, so if we consider only cell counts, we are only about half human. In terms of gene counts, the microbiome contains about 200 times more genes than the human genome, making bacterial genes responsible for over 99% of our body's gene content! Of all the microbial communities in the human body, the gut microbiome is by far the most dense, diverse, and physiologically important ecosystem to our overall health.

Relative Commensal Abundance	Result	Range	Units
 Bacteroidetes Phylum	71.280	50.000 - 95.000	%
 Firmicutes Phylum	20.040	3.500 - 40.000	%
 Proteobacteria Phylum	4.776	0.500 - 12.500	%
 Verrucomicrobia Phylum	3.903*H	0.000 - 2.400	%
 Actinobacteria Phylum	0.001	0.001 - 4.818	%
 Euryarchaeota Phylum	0.000	0.000 - 0.017	%

Your Phyla:



Healthy Phyla:



References:

NOTE: Relative abundance reference ranges have been based on a healthy population study.

King CH, et., al. (2019) Baseline human gut microbiota profile in healthy people and standard reporting template. PLoS One. 2019 Sep 11;14(9):e0206484.



3934192

Pathogen Summary:

Macroscopy Comment

BROWN coloured stool is considered normal in appearance.

Faecal Occult Blood Negative:

Faecal occult blood has not been detected in this specimen. If the test result is negative and clinical symptoms persist, additional follow-up testing using other clinical methods is recommended.

Metabolism Comment

In a healthy gut Short Chain Fatty Acids (SCFAs) exhibited in the following proportions; Butyrate, Acetate, Propionate (16%:60%:24%).

The primary SCFAs butyrate, propionate and acetate are produced by predominant commensal bacteria via fermentation of soluble dietary fibre and intestinal mucus glycans.

Key producers of SCFAs include *Faecalibacterium prausnitzii*, *Akkermansia muciniphila*, *Bacteroides fragilis*, *Bifidobacterium*, *Clostridium* and *Lactobacillus* Spp.

The SCFAs provide energy for intestinal cells and regulate the actions of specialised mucosal cells that produce anti-inflammatory and antimicrobial factors, mucins that constitute the mucus barriers, and gut active peptides that facilitate appetite regulation and euglycemia. Abnormal SCFAs may be associated with dysbiosis, intestinal barrier dysfunction and inflammatory conditions.

SCFA PRODUCTION TABLE

BACTERIA	BUTYRATE	PROPIONATE	ACETATE
<i>Akkermansia muciniphila</i>		▲▲▲	▲▲
<i>Anaerostipes caccae</i>	▲▲▲		
<i>Bacteroides</i> spp.		▲▲▲	
<i>Bifidobacterium</i> spp.		▲	▲▲▲
<i>Blautia obeum</i>		▲▲	▲▲▲
<i>Coprococcus eutactus</i>	▲		
<i>Escherichia coli</i>			▲
<i>Eubacterium rectale</i>	▲▲		
<i>Faecalibacterium prausnitzii</i>	▲▲▲		
<i>Lactobacillus</i> spp.	▲	▲	▲
<i>Roseburia homini</i>	▲▲		
<i>Ruminococcus bromii</i>	▲		▲
<i>Subdoligranulum variabile</i>	▲		

KEY

▲ Low Producers

▲▲ Moderate Producers

▲▲▲ High Producers

GIT Markers Comment

PANCREATIC ELASTASE: Normal exocrine pancreatic function.

Pancreatic Elastase reflects trypsin, chymotrypsin, amylase and lipase activity.

This test is not affected by supplements of pancreatic enzymes.

Healthy individuals should be producing >500 ug/g of PE-1 under normal/healthy conditions.

PE-1 levels between 200 - 500 ug/g may indicate suboptimal production.

PE-1 levels <200 ug/g indicate clear inadequate production.

The clinician should therefore consider digestive enzyme supplementation if one or more of the following conditions is present:

Loose watery stools, Undigested food in the stools, Post-prandial abdominal pain, Nausea or colicky abdominal pain, Gastroesophageal reflux symptoms, Bloating or food intolerance.

Testing performed by chemiluminescence immunoassay (CLIA).

ELEVATED CALPROTECTIN:

Elevated faecal calprotectin indicates a high probability of intestinal inflammation.



P: 1300 688 522
E: info@nutripath.com.au

-.GRETCHEN COOPER
70 VICTOR STREET
BANYO QLD 4014

JACQUI LEE
24-Sep-1983

Female

50 BARBER DRIVE
CAPALABA QLD 4157

LAB ID : 3934192
UR NO. : 6228912
Collection Date : 13-Nov-2023
Received Date: 15-Nov-2023



3934192

For patients with known inflammatory bowel disease in remission, faecal calprotectin above 50 ug/g is associated with an increased risk of relapse over the next 12 months. In patients with faecal calprotectin below 50u/g with strong clinical indications of intestinal inflammation, repeat testing may be useful. In small bowel Crohn's disease, the faecal calprotectin may not be elevated. Elevated faecal calprotectin may occur with other conditions including colorectal cancer, NSAID ulceration, coeliac disease, diverticulitis and chronic inflammation.

Further investigative procedures are necessary to determine the cause of inflammation.

Test performed by Phadia ELIA Fluorescence enzyme immunoassay (FEIA).

SECRETORY IGA NORMAL:

Secretory IgA is within range.

Secretory IgA represents the first line of defence of the gastrointestinal mucosa and is central to the normal function of the gastrointestinal tract as an immune barrier. Review this level with other pathogenic bacteria and normal commensal flora. Lower levels within reference range should be interpreted clinically.

ZONULIN NORMAL:

Zonulin is a protein that modulates intestinal barrier function. This results is considered normal.

BETA-GLUCURONIDASE NORMAL:

B-Glucuronidase is considered normal and is within reference range.

Opportunistic Bacteria Comment

METHANOBREVIBACTER SMITHII:

PHYLUM: Euryarchaeota

DESCRIPTION:

Methanobrevibacter smithii is a methane-producing microbe that plays an important role in the gut ecosystem by facilitating carbohydrate fermentation and production of short-chain fatty acids by commensal bacteria. Approximately 70% of the healthy cohort has below detectable levels of Methanobrevibacter smithii.

Phyla Microbiota Comment

VERRUCOMICROBIA (PHYLUM) ELEVATED:

DESCRIPTION:

Verrucomicrobia is a phylum of Gram-negative bacteria that contains only a few described species, found in the environment and gastrointestinal tract.

Akkermansia spp. is involved in gut membrane integrity and may be increased with polyphenols and prebiotics.

Verrucomicrobia aid in glucose homeostasis of the human gut and have anti-inflammatory properties that further aid in intestinal health.

TREATMENT SUGGESTIONS: Probiotic use and dietary modification use may assist in the rebalancing of microbial flora.



P: 1300 688 522
E: info@nutripath.com.au

-.GRETCHEN COOPER
70 VICTOR STREET
BANYO QLD 4014

JACQUI LEE
24-Sep-1983

Female

50 BARBER DRIVE
CAPALABA QLD 4157

LAB ID : 3934192
UR NO. : 6228912
Collection Date : 13-Nov-2023
Received Date: 15-Nov-2023



3934192

Normal Bacterial Flora Comment

BACTEROIDES FRAGILIS LOW:

PHYLUM: Bacteroidetes

DESCRIPTION: Bacteroides fragilis is an anaerobic, Gram-negative bacterium. It is part of the normal microbiota of the human colon and is generally commensal. Bacteroides fragilis plays an intricate role in the human colon and usually has a beneficial relationship with the host. Low Bacteroides fragilis levels have been associated with inflammatory bowel disease and Crohn's.

TREATMENT SUGGESTIONS: Treatment may involve the use of probiotics and dietary modification.

BIFIDOBACTERIUM SPECIES LOW:

PHYLUM: Actinobacteria

DESCRIPTION:

Bifidobacterium is a genus of gram-positive, nonmotile anaerobic bacteria that are ubiquitous inhabitants of the gastrointestinal tract and considered a probiotic. Bifidobacterium species prevent diarrhea and intestinal infections, alleviate constipation, and stimulate the immune system. Low levels may be associated with irritable bowel syndrome, asthma, autism, depressive disorder and with pathogenic bacterial infection.

TREATMENT SUGGESTIONS: Treatment may involve the use of Bifidobacterium containing probiotics and treatment of any intestinal infection.

BIFIDOBACTERIUM LONGUM LOW:

PHYLUM: Actinobacteria

DESCRIPTION:

Bifidobacterium longum is a Gram-positive, catalase-negative, rod-shaped bacterium present in the human gastrointestinal tract and one of the Bifidobacterium species. It can induce and regulate immune responses, reduce the expression of inflammatory cytokines, and maintain the normal intestinal barrier function.

Bifidobacterium longum is a clinically effective, well-established, multifunctional probiotic that has a long history of human use in alleviating gastrointestinal, immunological, and infectious diseases such as constipation, antibiotic associated diarrhoea, irritable bowel syndrome and ulcerative colitis. Low levels may be associated with irritable bowel syndrome, asthma, autism, depressive disorder and with pathogenic bacteria infection.

TREATMENT SUGGESTIONS: Treatment may involve the use of Bifidobacterium longum containing probiotics and treatment of any intestinal infections.

LACTOBACILLUS RHAMNOSUS LOW:

PHYLUM: Firmicutes

DESCRIPTION:

Lactobacillus Rhamnosus is a Gram-positive anaerobic bacterium and is one of the most widely used probiotic strains, of which various health effects are well documented including the prevention and treatment of gastro-intestinal infections and diarrhea and even preventing certain allergic symptoms.

Decreased Lactobacillus rhamnosus colonisation has been shown to decrease gastro-intestinal health, increasing the risk of gastro-intestinal infections and diarrhea as well as extra-intestinal infections including oral and respiratory health. Studies have also revealed that chronic psychological stress and alcohol use may be associated with a decrease in Lactobacillus species, as well as antibiotic / medication use.

TREATMENT SUGGESTIONS: Treatment may involve the use of Lactobacillus containing probiotics and treatment of any intestinal infections.

LACTOBACILLUS SPECIES LOW:

PHYLUM: Firmicutes

DESCRIPTION:

Lactobacillus species low. Lactobacillus is a genus of Gram-positive rod-shaped, non-spore-forming bacteria which constitute a



P: 1300 688 522
E: info@nutripath.com.au

-.GRETCHEN COOPER
70 VICTOR STREET
BANYO QLD 4014

JACQUI LEE
24-Sep-1983

Female

50 BARBER DRIVE
CAPALABA QLD 4157

LAB ID : 3934192
UR NO. : 6228912
Collection Date : 13-Nov-2023
Received Date: 15-Nov-2023



3934192

significantly important component of the human gastrointestinal system.

Lactobacillus exhibits a mutualistic relationship with the human body, as it protects the host against potential invasions by pathogens, and in turn, the host provides a source of nutrients. Lactobacilli are among the most common probiotic found in food such as yogurt, and it is diverse in its application to maintain gut health and to help treat ailments clinically such as diarrhea and obesity.

Lactobacillus species promote the anti-inflammatory response, thereby supporting the improvement of symptoms pertaining to asthma, chronic obstructive pulmonary disease, neuroinflammatory diseases, cardiovascular diseases, inflammatory bowel disease (IBD) and chronic infections in patients. Reduced levels increase the risk of infections and inflammation. Studies have also revealed that chronic psychological stress and alcohol use may be associated with a decrease in Lactobacillus species, as well as antibiotic / medication use.

TREATMENT SUGGESTIONS: Treatment may involve the use of Lactobacillus containing probiotics and treatment of any intestinal infections.

AKKERMANSIA MUCINIPHILA ELEVATED:

PHYLUM: Verrucomicrobia

DESCRIPTION:

Akkermansia muciniphila is a Gram-negative, strictly anaerobic, non-motile bacterium, often considered a human intestinal symbiont. There is growing evidence to suggest that the prevalence of this bacteria is associated with intestinal homeostasis, immunity, and a healthy gut. However, elevated colonisation may be associated with intestinal inflammation.

TREATMENT SUGGESTIONS: Treatment may involve the use of probiotics, treatment of any intestinal infections and dietary modification.



The Four “R” Treatment Protocol

REMOVE	Using a course of antimicrobial, antibacterial, antiviral or anti parasitic therapies in cases where organisms are present. It may also be necessary to remove offending foods, gluten, or medication that may be acting as antagonists. Consider testing IgG96 foods as a tool for removing offending foods.	ANTIMICROBIAL	Oil of oregano, berberine, caprylic acid
		ANTIBACTERIAL	Liquorice, zinc carnosine, mastic gum, tribulus, berberine, black walnut, caprylic acid, oil of oregano
		ANTIFUNGAL	Oil of oregano, caprylic acid, berberine, black walnut
		ANTIPARASITIC	Artemesia, black walnut, berberine, oil of oregano
		ANTIVIRAL	Cat's claw, berberine, echinacea, vitamin C, vitamin D3, zinc, reishi mushrooms
		BIOFILM	Oil of oregano, protease
REPLACE	In cases of maldigestion or malabsorption, it may be necessary to restore proper digestion by supplementing with digestive enzymes.	DIGESTIVE SUPPORT	Betaine hydrochloride, tilactase, amylase, lipase, protease, apple cider vinegar, herbal bitters
REINOCULATE	Recolonisation with healthy, beneficial bacteria. Supplementation with probiotics, along with the use of prebiotics helps re-establish the proper microbial balance.	PREBIOTICS	Slippery elm, pectin, larch arabinogalactans
		PROBIOTICS	Bifidobacterium animalis subsp. lactis, lactobacillus acidophilus, lactobacillus plantarum, lactobacillus casei, bifidobacterium breve, bifidobacterium bifidum, bifidobacterium longum, lactobacillus salivarius subsp. salivarius, lactobacillus paracasei, lactobacillus rhamnosus, Saccharomyces boulardii
REPAIR & REBALANCE	Restore the integrity of the gut mucosa by giving support to healthy mucosal cells, as well as immune support. Address whole body health and lifestyle factors so as to prevent future GI dysfunction.	INTESTINAL MUCOSA IMMUNE SUPPORT	Saccharomyces boulardii, lauric acid
		INTESTINAL BARRIER REPAIR	L-Glutamine, aloe vera, liquorice, marshmallow root, okra, quercetin, slippery elm, zinc carnosine, Saccharomyces boulardii, omega 3 essential fatty acids, B vitamins
		SUPPORT CONSIDERATION	Sleep, diet, exercise, and stress management