

Report to **IMEDICAL,**
1 Union st
, Pyrmont, NSW, 2009

Patient **HOWARTH, SOPHIE**
IMEDICAL C/O IMEDICAL 1 UNION ST
PYRMONT NSW 2009

Phone 0408403253
D.O.B 27/10/1970 Age 53 years Sex F

Ref. by/copy to IMEDICAL,
Collect date 17/06/2024 Lab ref 24-27878183
Collect time 08:34 AM Your ref
Reported 17/06/2024 07:33 PM

Tests requested SCP, LP1, FBE, IGE TPT*,FLO*

Clinical notes

SERUM HIGH SENSITIVITY C-REACTIVE PROTEIN (CRP)

Request Number 27878183
Date Collected 17 Jun 24
Time Collected 08:34
hsCRP (< 4.91) mg/L 0.48

The CDC / AHA recommend the following hsCRP cut-off points (tertiles) for cardiovascular disease risk assessment:

hsCRP level (mg/L)	Relative Risk
<1.0	Low
1.0 - 3.0	Average
>3.0	High

The average of two CRP tests, ideally taken two weeks apart, produces a more stable estimate of this marker.

A CRP greater than 10mg/L should prompt a search for a source of infection or inflammation.

**SURGERY
USE**

- Normal
- No Action/File
- Patient Notified
- Make Appoint.
- Further Tests
- Notes Required
- Speak with Dr.
- On Correct Treatment

Report to **IMEDICAL,**
1 Union st
, Pyrmont, NSW, 2009

Patient **HOWARTH, SOPHIE**
IMEDICAL C/O IMEDICAL 1 UNION ST
PYRMONT NSW 2009

Phone 0408403253
D.O.B 27/10/1970 Age 53 years Sex F

Ref. by/copy to IMEDICAL, Collect date 17/06/2024 Lab ref 24-27878183
Collect time 08:34 AM Your ref
Reported 17/06/2024 07:33 PM

Tests requested SCP, LP1, FBE, IGE TPT*,FLO*

Clinical notes

CELL DIFFERENTIATION MARKERS - LEUKAEMIA/LYMPHOCYTOSIS EVALUATION

POPULATION UNDER STUDY Peripheral Blood: All cells were analysed within the lymphoid scattergram.

Absolute Lymphocyte count 1.4 x10⁹ /L (1.0-4.0)

Antigen tested	Cell count %	Absolute count x10 ⁹ /L	Reference range x10 ⁹ /L
CD19 (B cell)	11	0.15	(0.04-0.50)
CD20 (B cell)	11	0.15	
CD23 (B cell subset)	2	0.03	
CD10 (CALLA +)	0	0.00	
CD5/CD19(B-CLL:NHL:Bsub)	< 1	< 0.01	
HLA-DR(Class II)	14	0.20	(0.04-0.50)
Kappa light chain	6	0.08	
Lambda light chain	5	0.07	
CD3 (Pan T cells)	83	1.16	(0.90-2.10)
CD4 (T helper)	46	0.64	(0.60-1.70)
CD5 (T cell/B sub)	82	1.15	
CD8(T suppressor)	33	0.46	(0.40-1.00)
Th:Ts(CD4/CD8)Ratio		1.39	(1.20-3.00)
CD3-/CD16+CD56+(NK)	6	0.08	(0.04-0.40)
CD3+/HLA-DR(act.T)	3	0.04	
CD13/CD33/CD34	<1		
CD117/CD34	0		

PHENOTYPE: Normal number of B-cells and these appear polyclonal and show no aberrant antigen expression. Normal number of T-cells and T-cell subsets with a normal CD4:CD8 ratio. Normal number of Natural Killer cells. No significant expression of the early stem marker CD34 or the myeloid markers CD13 or CD33.

CONCLUSION: No evidence of a clonal B-cell population or blast cell population in this study. Essentially normal peripheral blood lymphocyte subsets.

For queries contact Dr Lisa Clarke

SURGERY USE

Normal

No Action/File

Patient
Notified

Make
Appoint.

Further Tests

Notes
Required

Speak
with Dr.

On Correct
Treatment

