




# Microba *Insight*™

Gut microbiome report



**Name:** Christina Francis

**Sample ID:** BBT7682

**Report generated on:** 12-04-2024


# Introduction to *Microba Insight™* Report

Insight™ is a comprehensive report detailing key information about your personal gut microbiome. As new information about the links between the gut microbiome and health are revealed, we will continue to update your online report to include these new findings.

Any information provided by us (including any information contained on our website or in any microbiome report) is for information purposes only. Such information is not medical advice and must not be taken to be a substitute for a consultation with your healthcare professional or doctor. It is not intended to diagnose conditions nor prescribe the use of any remedy, diet or lifestyle practice. Your health is your responsibility and if you have any concerns related to your health we recommend that you seek the advice of your healthcare professional or doctor.

# Your report overview


Welcome to the start of your journey to understanding how your microbiome affects your health. Throughout this report, the analysed sample is compared to a healthy comparison group. This group is a collection of gut microbiome samples from everyday healthy people, who have not reported any significant health issues or symptoms. It represents a range of age groups, genders and diets.



## Microbial Diversity

### MICROBIAL DIVERSITY

Microbial diversity is a measure of the number of different microorganisms and the amount of each of these microorganisms in your sample. Average to high microbial diversity is associated with good health. A varied diet rich in plant-based foods such as fruits, vegetables, whole grains and nuts can help increase microbiome diversity. The Shannon Index is a measure of diversity which is used by members of the scientific community to compare results through time.



Your diversity level is

**Average**

Shannon Index

**4.15**

# Your microbial community

## YOUR TOP 5 MOST ABUNDANT SPECIES

Does your sample have an overly high abundance of any single organism? **No**

	Phylum	Species	Abundance	Range	Level
	Bacteroidota	<i>Bacteroides_A coprocola</i>	7.35%	0.00 - 0.156%	High
⊖	Proteobacteri a	<i>Escherichia coli</i>	5.99%	0.00 - 0.027%	High
⊕	Firmicutes_A	<i>Faecalibacterium prausnitzii_H</i>	5.14%	0.00 - 0.00%	High
⊕	Firmicutes_A	<i>Fusicatenibacter saccharivorans</i>	3.94%	0.518 - 7.34%	Average
⊕	Firmicutes_A	<i>Agathobacter rectale</i>	3.69%	0.00 - 7.22%	Average

# Your key insights

## Your gut microbiome's potential to produce **butyrate**, a primary fuel source for gut cells

**This is a good level!** Your potential to produce butyrate is at a level similar to the healthy group. To benefit from this important gut microbiome function, ensure your diet is rich in dietary sources of resistant starch.

A

This sample reported a level **similar to the healthy group**

The production of butyrate is a well-studied function of the gut microbiome. This 'metabolite' is named as one of the primary fuel sources for gut cells and has been shown to reduce inflammation throughout the body and help regulate appetite. A similar or high level to produce butyrate is beneficial for your gut microbiome and helps to maintain a healthy environment in the gut. Foods rich in resistant starch (e.g. green banana flour, raw oats or high amylose barley) will encourage microbes in your gut to produce butyrate.

EVIDENCE RATING ★★★★★

## Your microbiome's potential to contribute to **gut inflammation**

**This is not a good level.** Having a high potential to produce hexa-lipopolysaccharides (hexa-LPS) compared to the healthy group is not ideal. High levels of hexa-LPS can contribute to inflammation throughout the body. Avoiding excessive amounts of saturated fat could help to prevent this substance from spreading to areas outside of the gut. Also try adding foods high in fibre to reduce the levels of bacteria that produce this harmful substance.

A

This sample reported a level **higher than the healthy group**

Hexa-lipopolysaccharide (hexa-LPS) is a pro-inflammatory molecule and a component of the cell wall in some bacteria. When these bacteria die, hexa-LPS is released into the gut. Diets high in fat, especially saturated fat, allow hexa-LPS to cross the intestinal barrier and eventually enter the bloodstream. High levels of hexa-LPS in the blood have been observed in individuals with reduced heart health, variable glucose regulation, poor weight management, and poor liver health. If you have a high potential to produce hexa-LPS, you may wish to avoid excessive consumption of saturated fat. Dietary sources of saturated fat include butter, coconut oil, cheese, processed meats, chocolate, icecream, cakes and biscuits.

EVIDENCE RATING ★★★★★

# Your key insights

## Your gut microbiome's ability to break down fibre

**This is a good level!** Your potential to break down fibre is similar to the healthy group in this sample. This is an important gut microbiome function to maintain because it results in the production of beneficial substances that promote good gut health. To ensure the production of these beneficial compounds ensure your diet contains plenty of fibre.

**A****This sample reported a level similar to the healthy group**

Fibre-degrading bacteria are responsible for producing important by-products such as short chain fatty acids which play a critical role in keeping your gut healthy. Specific prebiotic fibres—detailed in your food suggestions—will promote the growth of your beneficial, fibre-degrading bacteria. A similar or high proportion of species that can break down fibre compared to the healthy group is considered beneficial.

EVIDENCE RATING ★★★★★

## Your gut microbiome's ability to break down protein

**This is a typical level.** The proportion of bacteria present in your sample that can break down protein is at level similar to the healthy group. When protein is broken down by bacteria in the gut microbiome it can lead to the production of substances that promote inflammation. To maintain this level, continue eating diverse sources of fibre to encourage the growth of your fibre-degrading bacteria instead of your protein-degrading bacteria.

**A****This sample reported a level similar to the healthy group**

Everyone's microbiome contains species that can break down protein into a variety of compounds, including some compounds that promote inflammation. Having a high proportion of these species may reflect an insufficient amount of fibre in the diet or an excessive intake of protein. A high proportion of protein-degrading bacteria suggests that not enough fibre is reaching the lower colon to feed the bacteria that specialise in eating fibre.

EVIDENCE RATING ★★★★★

# Your key insights

## Your microbiome's potential to produce branched chain amino acids

Your potential to produce branched chain amino acids is at a level similar to the healthy group. Similar or low levels of bacterially produced BCAAs compared to a healthy group is generally considered beneficial, as BCAAs are observed to be associated with poor weight management and variable glucose regulation.

**A****This sample reported a level similar to the healthy group**

BCAAs play an important role in building muscles and in helping regulate fat and sugar metabolism. However, a high potential to produce BCAAs by your gut microbiome may not be a good thing as high levels of bacterially produced BCAAs have been observed in individuals with poor weight management and variable glucose regulation. Having a low or similar potential to produce branched chain amino acids (BCAAs) compared to the healthy group is generally considered beneficial. Maintaining muscle mass through regular resistant exercise could help regulate BCAA blood levels.

EVIDENCE RATING ★★★★★

## Your microbiome's potential to influence and support your heart health

**This is a good level!** Your potential to produce trimethylamine (TMA) is at a level similar to the healthy group. Trimethylamine is converted by the human liver into trimethylamine oxide (TMAO) which has been linked to variable glucose regulation and reduced heart health. Plant compounds known as indoles have been shown to reduce the production of TMAO.

**A****This sample reported a level similar to the healthy group**

A similar or low potential to produce trimethylamine compared to the healthy group is generally considered beneficial. Trimethylamine is converted by the human liver into trimethylamine oxide (TMAO) which has been linked to poor heart and kidney health. Diets high in animal protein and low in fibre have been associated with increased trimethylamine production by gut microbes while plant chemicals known as indoles have been shown to reduce the production of TMAO.

EVIDENCE RATING ★★★★★



# Your key insights

## Your microbiome's potential to **protect your nervous system**

**This is a good level!** Your potential to produce indolepropionic acid (known as IPA) is similar to the healthy group. This is good, because IPA is a strong antioxidant that can protect nerve cells from damage and may help support glucose regulation.

**A****This sample reported a level **similar to the healthy group****

IPA is a strong antioxidant produced by our gut bacteria that performs many important functions in our gut. It can protect nerve cells from damage, suppress inflammation and may improve glucose regulation. Research suggests foods rich in ellagic acid (e.g. chestnuts and ellagic acid enriched pomegranate juice), as well as wholegrain wheat and rye may help support IPA production.

EVIDENCE RATING ★★☆☆☆

## Your microbiome's potential to **contribute to kidney health**

**This is a good level!** Your potential to degrade oxalates is similar to the healthy group. A similar or high potential to break down oxalate compared to the healthy group is generally considered beneficial.

**A****This sample reported a level **similar to the healthy group****

The gut microbiome of individuals who suffer from poor kidney health often have a low potential to degrade oxalate. Oxalate is one of the main influences of kidney health. If you have poor kidney health, you may need wish to discuss trialling a low oxalate diet with a health care professional. A similar or high potential to break down oxalate compared to the healthy group is generally considered beneficial.

EVIDENCE RATING ★★☆☆☆



# Your key insights

## Your gut microbiome's potential to produce strong-smelling flatulence

**This is not a good level.** Your potential to produce hydrogen sulphide is at a high level. Try adding foods high in prebiotic fibres, especially those foods containing resistant starch (RS) and fructooligosaccharides (FOS) such as cooked and cooled potatoes/sweet potatoes, rye, slightly green bananas, nectarines, and pears to reduce the production of hydrogen sulphide, which when produced by gut bacteria, has been associated with an impaired gut barrier function.

**A****This sample reported a level higher than the healthy group**

As the microbes in your gut digest different fuel sources, such as fibre, protein, mucus and even bile acids, they produce different types of gases as a by-product. Flatulence is primarily made up of odourless gases such as nitrogen, hydrogen, carbon dioxide, and methane. However, a small percent of flatulence can be made up of the gas hydrogen sulphide, which gives flatulence the characteristic rotten eggs smell. A small amount of hydrogen sulphide gas has been found to be protective of the gut, however a high potential to produce hydrogen sulphide has been associated with mitochondrial dysfunction and impaired gut barrier function. Research has found that the production of hydrogen sulphide by gut bacteria can be inhibited by consuming foods high in the prebiotic fibres resistant starch (RS) and fructooligosaccharides (FOS).

EVIDENCE RATING ★★☆☆☆

## Your microbiome's potential to contribute to cell replication and repair

**This is a good level!** Your gut microbiome's potential to produce folate is at a level similar to the healthy group. Folate is important for cell replication and repair. Your gut microbiome has the potential to contribute up to 37% of your daily folate requirement.


**A****This sample reported a level similar to the healthy group**

Folate plays an important role in cell replication and repair. Deficiencies can result in reduced heart and blood health. We cannot produce folate on our own and it is primarily obtained from plants in our diet (e.g. dark green leafy vegetables, fruits and legumes) and bacteria living in our gut. This bacterial production can supplement your body's folate requirements. A similar or high potential to produce folate compared to the healthy group is generally considered beneficial.


EVIDENCE RATING ★★☆☆☆

# Digging deeper into the detail

Gut microbiome report



# Sample Composition




# Microbiome Digestion Potential

The source of food that bacteria can use varies between different species of gut bacteria. Below we show the proportion of species in your gut microbiome that can break down the fuel sources fibre, protein and mucin (mucus). After you eat a meal, food gets broken down in your stomach and travels to your small intestine, where most nutrients are absorbed. The food components that cannot be absorbed in the small intestine, such as fibre and excess protein, make their way to your large intestine where your gut microbiota transform these components into a variety of products called metabolites. These metabolites can play an important role in your health. Read more about each of the fuel sources and their links to health on each of the tabs below.


## FIBRE

This scale indicates the proportion of species in your gut microbiome that can break down fibre. If you have a low proportion, consider adding more fibre to your diet to improve your gut health. Fibre is the main energy source of gut bacteria, who break it down into beneficial metabolites such as short chain fatty acids and B vitamins. Short chain fatty acids such as butyrate play an important role in keeping us healthy, and is one of the reasons fibre is an important component of a healthy diet.




## MUCIN

This scale indicates the proportion of species in your gut microbiome that can break down mucin, a component of the protective mucus layer that lines our gut. Some bacteria can use mucin as a fuel source. Mucus turnover is a normal part of our gut function, however when the abundance of bacteria that eat mucus becomes too high, this can result in a thinning of the mucus layer and activation of the immune system. Our mucus layer is important because it serves as a protective barrier between the cells lining our gut and harmful bacteria. Mucus-degrading bacteria may increase in abundance when there is not enough fibre reaching the lower large intestine, allowing gut bacteria that can use mucus as an energy source to multiply.



## PROTEIN

This scale indicates the proportion of species in your gut microbiome that can break down protein. If you have a high proportion, consider reducing the amount of protein in your diet to improve gut health. Although most protein is absorbed by your body, excess protein that is not absorbed will pass to your gut microbiome. The metabolites produced from the break down of protein are varied, with some being beneficial and others promoting inflammation. Diets high in animal protein and low in fibre have been observed to increase levels of pro-inflammatory gut metabolites.



# Microbial Metabolites

Your gut bacteria can produce thousands of different substances, called metabolites, when they use different fuel sources for energy. These metabolites can interact with our immune, metabolic and nervous systems to influence our health. Some of these metabolites promote good health while others promote poor health.

# Microbial Metabolites

Below we show the potential of your gut microbiome to produce or consume different metabolites associated with health. The microorganisms in your gut can transform the food components you eat into thousands of products called metabolites. Some metabolites have been associated with health benefits while others have been associated with poor health. Compare your microbiome's potential to produce and consume some of these metabolites with your selected comparison group. A ' + ' sign next to the compound name indicates it is associated with health benefits and a ' - ' sign indicates it is associated with poor health.

## HEALTH INDICATORS

### Produced

			ND	LOW	AVERAGE	HIGH
⊖	<b>Hexa-acylated lipopolysaccharide production</b>	<b>6.65%</b>				You
<b>The abundance of this metabolite is higher than the comparison group.</b>						
<p>Hexa-lipopolysaccharide (hexa-LPS) is a pro-inflammatory compound produced by some species of bacteria within the Proteobacteria phylum. High levels of hexa-LPS in the blood have been observed in individuals with metabolic and inflammatory conditions such as obesity, poor heart health, poor glucose regulation, and poor liver health. Diets high in fat, especially saturated fat, allow hexa LPS to cross the intestinal barrier and enter the bloodstream. Avoiding excessive intake of saturated fat can help reduce the ability of hexa-LPS to enter the bloodstream. Common dietary sources of saturated fats include butter, coconut products, palm oil, cheese, fatty meats, biscuit, cakes, chocolate and icecream.</p> <p><a href="#">[1]</a> <a href="#">[2]</a> <a href="#">[3]</a> <a href="#">[4]</a> <a href="#">[5]</a> <a href="#">[6]</a></p>						
			ND	LOW	AVERAGE	HIGH
⊖	<b>Methane production</b>	<b>0.781%</b>			You	
<b>The abundance of this metabolite is about the same as the comparison group.</b>						
<p>The gas methane can be produced by some species of the gut microbiome, primarily through the reduction of carbon dioxide and hydrogen. Although methane production is often detected in healthy adult populations, elevated levels of methane production has been associated with slower intestinal transit times and constipation.</p> <p><a href="#">[1]</a> <a href="#">[2]</a> <a href="#">[3]</a> <a href="#">[4]</a></p>						
			ND	LOW	AVERAGE	HIGH
⊖	<b>Trimethylamine production</b>	<b>2.87%</b>			You	
<b>The abundance of this metabolite is about the same as the comparison group.</b>						
<p>A high potential to produce trimethylamine has been correlated to poor heart health and poor glucose regulation. Once trimethylamine is produced by gut microbes, it is transported to the liver and converted to trimethylamine-n-oxide (TMAO). TMAO has been shown to be involved with blood sugar control, blood clotting and inflammation.</p> <p>The indoles diindolylmethane (DIM) and indole-3-carbinol (I3C) found in cruciferous vegetables (e.g. broccoli, cauliflower, cabbage, kale) may reduce the amount of trimethylamine that is converted to TMAO in the liver. In addition, excessive red meat consumption is associated with increased levels of TMAO in the blood. If your potential to produce trimethylamine is high, you may wish to increase your consumption of cruciferous vegetables and avoid eating excessive amounts of red meat. The Heart Foundation recommends limiting red meat consumption to less than 350g per week.</p> <p><a href="#">[1]</a> <a href="#">[2]</a> <a href="#">[3]</a> <a href="#">[4]</a> <a href="#">[5]</a> <a href="#">[6]</a> <a href="#">[7]</a></p>						

# Microbial Metabolites

HEALTH INDICATORS

## Produced

Ammonia (urease) production

7.95%

ND

LOW

AVERAGE

HIGH

You

**The abundance of this metabolite is about the same as the comparison group.**

Ammonia production is a normal way that bacteria recycle protein in the gut. However, high levels of ammonia production have been observed in individuals with impaired gut barrier function and inflammation of the gut.

[\[1\]](#) [\[2\]](#)

B. fragilis toxin production

0.00%

ND

LOW

AVERAGE

HIGH

You

**This metabolite is not detected in this microbiome.**

Most people's gut microbiome contain a species of bacteria called *Bacteroides fragilis*. A small proportion of *B. fragilis* strains have the ability to secrete a toxin. In some people this toxin can cause symptoms such as diarrhoea while other people can remain symptom free. There are concerns that this toxin can cause intestinal inflammation. If you are experiencing diarrhoea and have this toxin, consider seeing a healthcare practitioner.

[\[1\]](#) [\[2\]](#)

Beta-glucuronidase production

27.6%

ND

LOW

AVERAGE

HIGH

You

**The abundance of this metabolite is higher than the comparison group.**

Beta-glucuronidase is a bacterial enzyme which can limit the excretion of compounds from the body such as medications, hormones and environmental toxins. One human study has suggested that consuming glucomannan can reduce faecal beta-glucuronidase activity. Glucomannan is a type of prebiotic fibre found in konjac root which is commonly used to make low calorie pasta and noodles.

[\[1\]](#) [\[2\]](#)

Hydrogen sulphide production

16.7%

ND

LOW

AVERAGE

HIGH

You

**The abundance of this metabolite is higher than the comparison group.**

The gas hydrogen sulphide is produced by bacteria when they break down sulphur-containing amino acids found in foods such as eggs, meat, and fish. This gas is responsible for the rotten egg smell of flatulence. At low to average levels, hydrogen sulphide can play a beneficial role by acting as an energy source for gut cells. However at high levels hydrogen sulphide can inhibit energy production in gut cells and disrupt the gut mucus barrier. Elevated levels of hydrogen sulphide have been associated with poor intestinal health. Laboratory based studies have suggested that eating foods high in resistant starch (e.g. lentils, peas, beans, rolled oats and cooked and cooled potatoes) or fructooligosaccharides (FOS) (e.g. onions, garlic, leek, banana, peaches, wheat, barley) can reduce the production of hydrogen sulphide by the microbiome.

[\[1\]](#) [\[2\]](#)

15/71 • v0.12.3-gb81b11 • MGDB\_v2 • BBT7682 • Christina Francis • 12/04/2024

Any information provided by us (including any information contained on our website or in any microbiome report) is for information purposes only. Such information is not medical advice and must not be taken to be a substitute for a consultation with your health care professional or doctor. It is not intended to diagnose conditions nor prescribe the use of any remedy, diet or lifestyle practice. Your health is your responsibility and if you have any concerns related to your health we recommend that you seek the advice of your health care professional or doctor.






# Microbial Metabolites

HEALTH INDICATORS

## Consumed



## Produced



[1] [2] [3]



[1] [2] [3] [4] [5]




[1] [2] [3]

# Microbial Metabolites

## NEUROENDOCRINE

## Consumed



**The abundance of this metabolite is higher than the comparison group.**

GABA (gamma-aminobutyric acid) plays an important role in regulating mental state by calming the nervous system. Low levels of GABA being associated with poor mental health. Most GABA is produced in the brain however your gut microbiome may contribute to your GABA levels as some bacteria can produce or consume GABA. The role of gut bacteria that produce GABA in mental health is currently not well understood. If you are concerned about your mental health, it is important to seek professional help.

[1] [2] [3]

## Produced



[1] [2] [3] [4]



[1] [2]




[1] [2] [3] [4] [5] [6]

# Microbial Metabolites

## SHORT CHAIN FATTY ACIDS

## Produced



**The abundance of this metabolite is about the same as the comparison group.**

Acetate is the most abundant short chain fatty acid produced by our gut microbiome. It plays a beneficial role by suppressing inflammation, regulating appetite, and regulating fat metabolism. Several bacterial species can also convert acetate to the beneficial short chain fatty acid, butyrate. Laboratory studies have shown that consuming pectin can increase acetate production. Pectin rich foods include avocado, berries, kiwi, pumpkin and zucchini.

[\[1\]](#) [\[2\]](#) [\[3\]](#) [\[4\]](#)

## Produced





## Produced



# Species of Interest

## BACTERIA (PROKARYOTES)

### Agathobacter

DETECTED




*Agathobacter rectale*

You

3.69%

Average

2.43%




*Agathobacter faecis*

You

0.875%

Average

1.39%




*Agathobacter*  
*sp000434275*

You

0.202%

Average

0.038%




*Agathobacter*  
*sp900317585*

You

0.038%

Average

0.005%



### Akkermansia

NOT DETECTED

### Bifidobacterium

DETECTED




*Bifidobacterium animalis*

You

0.226%

Average

0.089%



Bilophila


DETECTED



Bilophila wadsworthia

You0.022%

Average0.081%



Campylobacter

NOT DETECTED

Citrobacter

NOT DETECTED

Clostridioides

NOT DETECTED


Clostridium

DETECTED

Clostridium\_M  
sp000431375

You0.030%


Average0.100%



Clostridium\_Q  
sp003024715

You0.066%

Average0.116%



Corynebacterium

NOT DETECTED

Desulfovibrio

NOT DETECTED

Eggerthella

NOT DETECTED

Enterobacter

NOT DETECTED

Escherichia


DETECTED



Escherichia coli

You5.99%

Average0.093%



Faecalibacterium


DETECTED

⊕

**Faecalibacterium  
prausnitzii\_H**

You  
Average


**5.14%**  
**0.103%**



**Faecalibacterium  
prausnitzii\_D**

You  
Average

**0.642%**  
**0.705%**




⊕

**Faecalibacterium  
prausnitzii\_C**

You  
Average


**0.597%**  
**0.926%**



**Faecalibacterium  
prausnitzii\_G**

You  
Average


**0.341%**  
**1.03%**



**Faecalibacterium  
prausnitzii\_I**

You  
Average

**0.283%**  
**0.236%**




⊕

**Faecalibacterium  
prausnitzii\_K**

You  
Average


**0.139%**  
**0.644%**



**Faecalibacterium  
MIC7145**

You  
Average


**0.102%**  
**0.662%**



**Faecalibacterium  
MIC8666**

You  
Average


**0.092%**  
**0.260%**



*Faecalibacterium  
prausnitzii\_A*

You  
Average

0.091%  
0.114%



*Fusobacterium*

NOT DETECTED

*Helicobacter*

NOT DETECTED

*Klebsiella*


DETECTED



*Klebsiella pneumoniae*

You  
Average

0.665%  
0.004%



*Lactobacillus*


DETECTED



*Lactobacillus\_C  
rhamnosus*

You  
Average

0.010%  
0.018%



*Oxalobacter*

NOT DETECTED

*Porphyromonas*

NOT DETECTED

*Prevotella*

NOT DETECTED

*Roseburia*


DETECTED



*Roseburia intestinalis*

You  
Average


1.05%  
0.457%



*Roseburia hominis*

You  
Average

0.198%  
0.119%




Ruminococcus

DETECTED


Ruminococcus\_A  
sp003011855

You 0.358%  
Average 0.364%




Ruminococcus\_A  
sp000437095

You 0.096%  
Average 0.069%




Ruminococcus\_D  
bicirculans

You 0.218%  
Average 1.18%



Ruminococcus\_E  
bromii\_B

You 1.76%  
Average 2.00%



Salmonella

NOT DETECTED


Streptococcus

DETECTED




Streptococcus  
thermophilus

You 0.473%  
Average 0.080%



Streptococcus salivarius

You 0.045%  
Average 0.109%




# Species of Interest

## ARCHAEA (PROKARYOTES)

### Methanogens

DETECTED



### Other Archea

NOT DETECTED




# Species of Interest

## YEASTS/FUNGI & PROTISTS (EUKARYOTES)

*Blastocystis*

DETECTED



*Candida*

NOT DETECTED

*Saccharomyces*


NOT DETECTED

*Other Eukaryotes*

NOT DETECTED


# Microbial Profile

This section shows the different bacteria, archaea and eukaryotes present in your gut. A phylum is the highest level of grouping (comprising hundreds to thousands of species), whereas a species is the most detailed view of your gut microbiome



# Your Microbiome Profile

PHYLUM



This Sample

Phylum		Abundance	Range	Level
	Firmicutes_A	60.9%	41.8 - 73.1%	Average
	Bacteroidota	15.7%	10.4 - 26.4%	Average
	Proteobacteria	7.30%	0.315 - 3.02%	High
	Firmicutes	2.03%	0.741 - 9.75%	Average
	Euryarchaeota	0.809%	0.00 - 0.717%	High
	Firmicutes_C	0.683%	0.271 - 1.38%	Average
	Actinobacteriota	0.528%	1.03 - 7.36%	Low
	Desulfobacterota_A	0.042%	0.020 - 0.385%	Average
	Eukaryote_unclassified	0.035%	0.00 - 0.00%	High

# Species Profile

SPECIES

Phylum	Species	Abundance	Range	Level
Bacteroidota	<i>Bacteroides_A coprocola</i>	7.35%	0.00 - 0.156%	High
<p>This is a common inhabitant of the gut.</p> <p><b>Fuel Sources Used:</b> This species is a good degrader of fibre, a good degrader of mucin, and a moderate degrader of protein.</p> <p><b>Metabolites produced:</b> Our genomic analysis indicates that most members of this species can produce the following metabolites: acetate, biotin (B7), branched chain amino acids, folate (B9), lactate, riboflavin (B2), vitamin K.</p> <p><b>Metabolites consumed:</b> In addition, the genomic analysis indicates that most members of this species do not consume any reported metabolites.</p> <p><b>Emerging Research:</b> This species has multiple strains, with likely multiple roles. One study observed different strains of this species in patients with type 2 diabetes compared to healthy controls. Another study showed different strains of this species were present in individuals in Europe compared to the USA.</p>				
⊖ Proteobacteria	<i>Escherichia coli</i>	5.99%	0.00 - 0.027%	High
<p>This species is a common inhabitant of the gut, although it is usually present at a low abundance compared to other gut microbiome species.</p> <p><b>Fuel Sources Used:</b> This species is a moderate degrader of fibre, a poor degrader of mucin, and a moderate degrader of protein.</p> <p><b>Metabolites produced:</b> Our genomic analysis indicates that most members of this species can produce the following metabolites: acetate, beta-glucuronidase, biotin (B7), branched chain amino acids, folate (B9), GABA, hexa-LPS, hydrogen sulphide, lactate, propionate, riboflavin (B2), vitamin K.</p> <p><b>Metabolites consumed:</b> In addition, the genomic analysis shows that most members of this species can consume: GABA, oxalate.</p> <p><b>Emerging Research:</b> This species encompasses a large number of strains with diverse properties; a few well-known strains are a common cause of poor intestinal health. However, most strains will not cause health problems. Studies have observed this species at higher levels in individuals with poor intestinal and liver health. Additionally, a recent study identified several strains from this species can produce a toxin called colibactin which can lead to inflammation and damage DNA in human cells.</p>				

# Species Profile

## SPECIES

	Phylum	Species	Abundance	Range	Level
+	Firmicutes_A	<i>Faecalibacterium prausnitzii_H</i>	5.14%	0.00 - 0.00%	High
<p>This is an important member of the gut microbiome.</p> <p><b>Fuel Sources Used:</b> This species is a moderate degrader of fibre, a poor degrader of mucin, and a moderate degrader of protein.</p> <p><b>Metabolites produced:</b> Our genomic analysis indicates that most members of this species can produce the following metabolites: acetate, beta-glucuronidase, branched chain amino acids.</p> <p><b>Metabolites consumed:</b> In addition, the genomic analysis indicates that most members of this species do not consume any reported metabolites.</p> <p><b>Emerging Research:</b> This species can also use branched chain amino acids and the short chain fatty acid acetate for energy.</p> <p>This species has recently been split into several different species, and low levels of some of these species have been observed in individuals with poor intestinal health, ulcerative colitis, colon cancer, type II diabetes and chronic fatigue syndrome.</p>					
+	Firmicutes_A	<i>Fusicatenibacter saccharivorans</i>	3.94%	0.518 - 7.34%	Average
<p>This is a recently discovered species and an inhabitant of the human gut.</p> <p><b>Fuel Sources Used:</b> This species is a moderate degrader of fibre, mucin, and protein.</p> <p><b>Metabolites produced:</b> Our genomic analysis indicates that most members of this species can produce the following metabolites: acetate, ammonia (urease), beta-glucuronidase, branched chain amino acids, cobalamin (B12), folate (B9), hydrogen sulphide, lactate.</p> <p><b>Metabolites consumed:</b> In addition, the genomic analysis indicates that most members of this species do not consume any reported metabolites.</p> <p><b>Emerging Research:</b> Lower levels of this species were observed in people with poor intestinal health.</p>					

# Species Profile

## SPECIES

Phylum	Species	Abundance	Range	Level
<div><div>+</div><div>Firmicutes_A</div></div>	<i>Agathobacter rectale</i>	3.69%	0.00 - 7.22%	Average
<p>Previously named <i>Eubacterium rectale</i>, this is a common member of the human gut.</p> <p><b>Fuel Sources Used:</b> This species is a moderate degrader of fibre, a poor degrader of mucin, and a moderate degrader of protein.</p> <p><b>Metabolites produced:</b> Our genomic analysis indicates that most members of this species can produce the following metabolites: acetate, branched chain amino acids, butyrate, cobalamin (B12), folate (B9), lactate, riboflavin (B2).</p> <p><b>Metabolites consumed:</b> In addition, the genomic analysis indicates that most members of this species do not consume any reported metabolites.</p> <p><b>Emerging Research:</b> This species is commonly associated with healthier people in research studies. Lower levels of this bacterium have been observed in people with poor intestinal health.</p>				
<div><div></div><div>Firmicutes_A</div></div>	<i>GCA-900066995 sp900291955</i>	3.68%	0.00 - 0.00%	High
<p><b>Fuel Sources Used:</b> This species is a poor degrader of fibre, a poor degrader of mucin, and a moderate degrader of protein.</p> <p><b>Metabolites produced:</b> Our genomic analysis indicates that most members of this species can produce the following metabolites: acetate, branched chain amino acids, lactate, riboflavin (B2).</p> <p><b>Metabolites consumed:</b> In addition, the genomic analysis indicates that most members of this species do not consume any reported metabolites.</p> <p><b>Emerging Research:</b> This species is named according to the Genome Taxonomy Database (GTDB), a standardised microbial taxonomy based on the genomes of organisms, rather than their phenotype.</p>				

# Species Profile

SPECIES

Phylum	Species	Abundance	Range	Level
Firmicutes_A	<i>UBA11524 sp000437595</i>	2.51%	0.00 - 2.87%	Average
<p><b>Fuel Sources Used:</b> This species is a moderate degrader of fibre, a poor degrader of mucin, and a moderate degrader of protein.</p> <p><b>Metabolites produced:</b> Our genomic analysis indicates that most members of this species can produce the following metabolites: acetate, beta-glucuronidase, branched chain amino acids.</p> <p><b>Metabolites consumed:</b> In addition, the genomic analysis indicates that most members of this species do not consume any reported metabolites.</p> <p><b>Emerging Research:</b> This species is named according to the Genome Taxonomy Database (GTDB), a standardised microbial taxonomy based on the genomes of organisms, rather than their phenotype.</p>				
Firmicutes_A	<i>Blautia_A wexlerae</i>	2.24%	0.344 - 6.60%	Average
<p>This is a recently discovered and common inhabitant of the gut.</p> <p><b>Fuel Sources Used:</b> This species is a moderate degrader of fibre, a poor degrader of mucin, and a moderate degrader of protein.</p> <p><b>Metabolites produced:</b> Our genomic analysis indicates that most members of this species can produce the following metabolites: acetate, ammonia (urease), biotin (B7), branched chain amino acids, cobalamin (B12), folate (B9), lactate, riboflavin (B2).</p> <p><b>Metabolites consumed:</b> In addition, the genomic analysis indicates that most members of this species do not consume any reported metabolites.</p>				



# Species Profile

## SPECIES

Phylum	Species	Abundance	Range	Level	
+	Firmicutes_A	<i>Ruminiclostridium_E siraeum</i>	2.15%	0.00 - 1.58%	High
<p>Formerly known as <i>Eubacterium siraeum</i>. This is a common inhabitant of the human gut.</p> <p><b>Fuel Sources Used:</b> This species is a moderate degrader of fibre, a good degrader of mucin, and a moderate degrader of protein.</p> <p><b>Metabolites produced:</b> Our genomic analysis indicates that most members of this species can produce the following metabolites: acetate, branched chain amino acids, folate (B9).</p> <p><b>Metabolites consumed:</b> In addition, the genomic analysis indicates that most members of this species do not consume any reported metabolites.</p> <p><b>Emerging Research:</b> This species was observed at lower levels in individuals with poor metabolic and intestinal health, suggesting it likely plays a role in promoting good health.</p>					
	Firmicutes_A	<i>Blautia_A obeum</i>	2.07%	0.00 - 1.41%	High
<p><b>Fuel Sources Used:</b> This species is a moderate degrader of fibre, a poor degrader of mucin, and a moderate degrader of protein.</p> <p><b>Metabolites produced:</b> Our genomic analysis indicates that most members of this species can produce the following metabolites: acetate, biotin (B7), branched chain amino acids, cobalamin (B12), folate (B9), lactate, propionate.</p> <p><b>Metabolites consumed:</b> In addition, the genomic analysis indicates that most members of this species do not consume any reported metabolites.</p>					


# Species Profile

SPECIES

Phylum	Species	Abundance	Range	Level
Firmicutes_A	<i>Ruminococcus_E bromii_B</i>	1.76%	0.00 - 6.16%	Average
⊕ Firmicutes_A	<i>Coprococcus eutactus_A</i>	1.73%	0.00 - 1.43%	High
<p>This is a common inhabitant of the human gut.</p> <p><b>Fuel Sources Used:</b> This species is a moderate degrader of fibre, mucin, and protein.</p> <p><b>Metabolites produced:</b> Our genomic analysis indicates that most members of this species can produce the following metabolites: acetate, branched chain amino acids, butyrate, folate (B9), lactate, riboflavin (B2).</p> <p><b>Metabolites consumed:</b> In addition, the genomic analysis indicates that most members of this species do not consume any reported metabolites.</p> <p><b>Emerging Research:</b> This species has been observed at lower levels in individuals with lung conditions and poor intestinal health, and children with poor glucose regulation, indicating it likely plays a beneficial role in health.</p>				
Firmicutes_A	<i>Gemmiger sp003476825</i>	1.68%	0.00 - 2.86%	Average
Bacteroidota	<i>Bacteroides stercoris</i>	1.63%	0.00 - 2.66%	Average
Firmicutes_A	<i>Lachnospira eligens_B</i>	1.57%	0.00 - 1.74%	Average
Firmicutes_A	<i>Blautia_A sp900066165</i>	1.57%	0.193 - 2.41%	Average
Firmicutes_A	<i>Coprococcus sp900066115</i>	1.55%	0.00 - 0.867%	High

# Species Profile

SPECIES

	Phylum	Species	Abundance	Range	Level
	Bacteroidota	<i>Bacteroides uniformis</i>	1.37%	0.136 - 4.89%	Average
<p>This is one of the most common inhabitants of the human gut.</p> <p><b>Fuel Sources Used:</b> This species is a good degrader of fibre, a good degrader of mucin, and a moderate degrader of protein.</p> <p><b>Metabolites produced:</b> Our genomic analysis indicates that most members of this species can produce the following metabolites: acetate, beta-glucuronidase, biotin (B7), branched chain amino acids, folate (B9), GABA, lactate, riboflavin (B2).</p> <p><b>Metabolites consumed:</b> In addition, the genomic analysis indicates that most members of this species do not consume any reported metabolites.</p> <p><b>Emerging Research:</b> Certain strains have been observed to promote the production of anti-inflammatory compounds, and improve health in mouse models, however this has not yet been validated in humans. One study observed higher levels of this species in patients with poor intestinal health.</p>					
	Firmicutes_A	<i>KLE1615 sp900066985</i>	1.25%	0.172 - 2.22%	Average
	Firmicutes_A	<i>UBA1394 sp900066845</i>	1.23%	0.00 - 0.426%	High

# Species Profile

SPECIES

	Phylum	Species	Abundance	Range	Level
⊖	Bacteroidota	<i>Bacteroides_B vulgatus</i>	1.22%	0.00 - 6.38%	Average
<p>This is one of the most common inhabitants of the human gut.</p> <p><b>Fuel Sources Used:</b> This species is a good degrader of fibre, a good degrader of mucin, and a moderate degrader of protein.</p> <p><b>Metabolites produced:</b> Our genomic analysis indicates that most members of this species can produce the following metabolites: acetate, biotin (B7), branched chain amino acids, folate (B9), GABA, lactate, riboflavin (B2), vitamin K.</p> <p><b>Metabolites consumed:</b> In addition, the genomic analysis shows that most members of this species can consume: GABA.</p> <p><b>Emerging Research:</b> Higher levels of this bacteria have been associated with a wide range of poor health conditions. This species has been associated with a diet high in red meat.</p>					
	Firmicutes_A	<i>TF01-11 sp001414325</i>	1.19%	0.00 - 0.475%	High
	Bacteroidota	<i>CAG-485 MIC8356</i>	1.07%	0.00 - 0.00%	High
	Firmicutes_A	<i>CAG-217 sp000436335</i>	1.06%	0.00 - 1.30%	Average



# Species Profile

## SPECIES

	Phylum	Species	Abundance	Range	Level
+	Firmicutes_A	<i>Roseburia intestinalis</i>	1.05%	0.00 - 1.12%	Average
<p>This is a common and important member of the human gut microbiome.</p> <p><b>Fuel Sources Used:</b> This species is a good degrader of fibre, a moderate degrader of mucin, and a moderate degrader of protein.</p> <p><b>Metabolites produced:</b> Our genomic analysis indicates that most members of this species can produce the following metabolites: acetate, ammonia (urease), beta-glucuronidase, biotin (B7), branched chain amino acids, butyrate, cobalamin (B12), folate (B9), hydrogen sulphide, lactate, riboflavin (B2).</p> <p><b>Metabolites consumed:</b> In addition, the genomic analysis indicates that most members of this species do not consume any reported metabolites.</p> <p><b>Emerging Research:</b> This is a primary producer of the beneficial short chain fatty acid butyrate. This species has been observed at lower levels in individuals with poor glucose regulation, as well as younger people with poor intestinal health, suggesting this species likely plays a beneficial role in health.</p>					
-	Bacteroidota	<i>Bacteroides_B massiliensis</i>	0.921%	0.00 - 1.39%	Average
<p>This is a normal inhabitant of the human gut microbiome.</p> <p><b>Fuel Sources Used:</b> This species is a moderate degrader of fibre, a good degrader of mucin, and a moderate degrader of protein.</p> <p><b>Metabolites produced:</b> Our genomic analysis indicates that most members of this species can produce the following metabolites: acetate, biotin (B7), branched chain amino acids, folate (B9), GABA, lactate, riboflavin (B2), vitamin K.</p> <p><b>Metabolites consumed:</b> In addition, the genomic analysis indicates that most members of this species do not consume any reported metabolites.</p> <p><b>Emerging Research:</b> Higher levels of this species have been observed in individuals with poor intestinal health. This species has also been associated with a diet high in red meat.</p>					

# Species Profile

## SPECIES

Phylum	Species	Abundance	Range	Level
 Firmicutes_A	<i>Dorea longicatena</i>	0.899%	0.00 - 1.76%	Average
<p>This is a common inhabitant of the human gut.</p> <p><b>Fuel Sources Used:</b> This species is a moderate degrader of fibre, a poor degrader of mucin, and a moderate degrader of protein.</p> <p><b>Metabolites produced:</b> Our genomic analysis indicates that most members of this species can produce the following metabolites: acetate, branched chain amino acids, cobalamin (B12), folate (B9), lactate, riboflavin (B2), trimethylamine.</p> <p><b>Metabolites consumed:</b> In addition, the genomic analysis indicates that most members of this species do not consume any reported metabolites.</p> <p><b>Emerging Research:</b> Multiple studies have observed this species at higher levels in individuals with obesity compared to lean controls. However in obese, postmenopausal women, this species was associated with improved insulin sensitivity. This species has also been observed at lower levels in individuals with chronic fatigue syndrome and poor intestinal health.</p>				
 Firmicutes_A	<i>Agathobacter faecis</i>	0.875%	0.00 - 4.17%	Average
<p>Previously named <i>Roseburia faecis</i>. This is a common and important member of the human gut microbiome.</p> <p><b>Fuel Sources Used:</b> This species is a moderate degrader of fibre, mucin, and protein.</p> <p><b>Metabolites produced:</b> Our genomic analysis indicates that most members of this species can produce the following metabolites: acetate, branched chain amino acids, butyrate, folate (B9), lactate, riboflavin (B2).</p> <p><b>Metabolites consumed:</b> In addition, the genomic analysis indicates that most members of this species do not consume any reported metabolites.</p> <p><b>Emerging Research:</b> This species is commonly associated with healthy people in research studies. Low levels of this bacterium have been observed in patients with poor liver health.</p>				

# Species Profile

## SPECIES

	Phylum	Species	Abundance	Range	Level
	Firmicutes_A	<i>Blautia_A massiliensis</i>	0.746%	0.00 - 1.83%	Average
	Firmicutes_C	<i>Dialister succinatiphilus</i>	0.683%	0.00 - 0.00%	High
⊖	Proteobacteria	<i>Klebsiella pneumoniae</i>	0.665%	0.00 - 0.00%	High

This species can be found in several body sites, such as the mouth, nose, lungs and gut.

### Fuel Sources Used:

This species is a moderate degrader of fibre, a poor degrader of mucin, and a moderate degrader of protein.

### Metabolites produced:

Our genomic analysis indicates that most members of this species can produce the following metabolites: acetate, ammonia (urease), biotin (B7), branched chain amino acids, cobalamin (B12), folate (B9), GABA, hexa-LPS, hydrogen sulphide, lactate, propionate, riboflavin (B2), trimethylamine, vitamin K.

### Metabolites consumed:

In addition, the genomic analysis shows that most members of this species can consume: GABA.

### Emerging Research:

Although this species is present in healthy individuals, an overgrowth of this species in the gut has been associated with poor heart health, poor intestinal health, and poor liver health.

	Firmicutes_A	<i>Lachnospira rogosae</i>	0.645%	0.00 - 1.05%	Average
	Firmicutes_A	<i>Faecalibacterium prausnitzii_D</i>	0.642%	0.00 - 1.72%	Average
	Firmicutes_A	<i>CAG-41 sp900066215</i>	0.637%	0.00 - 0.815%	Average
	Firmicutes	<i>CAG-302 MIC8491</i>	0.623%	0.00 - 0.016%	High

# Species Profile

## SPECIES

	Phylum	Species	Abundance	Range	Level
+	Euryarchaeota	<i>Methanobrevibacter_A smithii</i>	0.600%	0.00 - 0.583%	High
<p>This is a single celled organism belonging to the Archaea domain and is the most common archaeal species found in the human gut.</p> <p><b>Fuel Sources Used:</b> This species is a poor degrader of fibre, a poor degrader of mucin, and a moderate degrader of protein.</p> <p><b>Metabolites produced:</b> Our genomic analysis indicates that most members of this species can produce the following metabolites: acetate, branched chain amino acids, histamine, methane.</p> <p><b>Metabolites consumed:</b> In addition, the genomic analysis indicates that most members of this species do not consume any reported metabolites.</p> <p><b>Emerging Research:</b> Higher levels of <i>M. smithii</i> have been observed with constipation, diverticulosis, and other conditions. However, lower levels have been observed in patients with poor intestinal health. <i>M. smithii</i> plays an important role in the gut because it is one of the few species that can remove excess hydrogen.</p>					
+	Firmicutes_A	<i>Faecalibacterium prausnitzii_C</i>	0.597%	0.00 - 2.02%	Average
<p><i>Faecalibacterium prausnitzii_C</i> (aka strain A2-165) is an important member of the human gut microbiome.</p> <p><b>Fuel Sources Used:</b> This species is a moderate degrader of fibre, a poor degrader of mucin, and a moderate degrader of protein.</p> <p><b>Metabolites produced:</b> Our genomic analysis indicates that most members of this species can produce the following metabolites: acetate, beta-glucuronidase, branched chain amino acids.</p> <p><b>Metabolites consumed:</b> In addition, the genomic analysis indicates that most members of this species do not consume any reported metabolites.</p> <p><b>Emerging Research:</b> Low levels of <i>F. prausnitzii</i> have been linked to a range of poor health conditions, including intestinal, metabolic, and mental health.</p>					



# Species Profile

## SPECIES

	Phylum	Species	Abundance	Range	Level
+	Firmicutes_A	<i>Eubacterium_E hallii</i>	0.582%	0.00 - 1.75%	Average
<p>This is an important member of the gut microbiome.</p> <p><b>Fuel Sources Used:</b> This species is a poor degrader of fibre, a poor degrader of mucin, and a moderate degrader of protein.</p> <p><b>Metabolites produced:</b> Our genomic analysis indicates that most members of this species can produce the following metabolites: acetate, branched chain amino acids, butyrate, cobalamin (B12), folate (B9), lactate, riboflavin (B2).</p> <p><b>Metabolites consumed:</b> In addition, the genomic analysis indicates that most members of this species do not consume any reported metabolites.</p> <p><b>Emerging Research:</b> This species can also use the short chain fatty acid acetate, as well as lactate produced by bacterial species such as <i>Bifidobacterium spp.</i> for energy.</p> <p>One study observed lower levels of this species in people with poor intestinal health. Also, a study in mice suggested this species may be effective at improving insulin sensitivity.</p>					
	Bacteroidota	<i>Bacteroides xylanisolvens</i>	0.549%	0.00 - 0.429%	High
	Firmicutes_A	<i>Gemmiger formicilis</i>	0.545%	0.00 - 2.51%	Average
	Firmicutes_A	<i>Anaerostipes hadrus_A</i>	0.508%	0.00 - 0.655%	Average


# Species Profile

## SPECIES

	Phylum	Species	Abundance	Range	Level
+	Firmicutes	<i>Streptococcus thermophilus</i>	0.473%	0.00 - 0.221%	High
<p>This is the most widely used lactate bacteria in the dairy industry for producing cheese and yogurt (it is considered the safest <i>Streptococcus</i> species by the dairy industry). It helps make reduced-fat cheese with similar characteristics to full-fat cheese.</p> <p><b>Fuel Sources Used:</b> This species is a poor degrader of fibre, a poor degrader of mucin, and a moderate degrader of protein.</p> <p><b>Metabolites produced:</b> Our genomic analysis indicates that most members of this species can produce the following metabolites: acetate, ammonia (urease), branched chain amino acids, folate (B9), lactate.</p> <p><b>Metabolites consumed:</b> In addition, the genomic analysis indicates that most members of this species do not consume any reported metabolites.</p> <p><b>Emerging Research:</b> Because of its superior ability to use lactose, this species is often used by lactose-intolerant individuals to help them digest milk products. This bacterium also shows good potential for reducing inflammation, although more research needs to be conducted in humans.</p>					
+	Firmicutes_A	<i>Faecalicatena lactaris</i>	0.440%	0.00 - 0.879%	Average
<p>Previously called <i>Ruminococcus lactaris</i>, this is a common inhabitant of the human gut.</p> <p><b>Fuel Sources Used:</b> This species is a poor degrader of fibre, a moderate degrader of mucin, and a moderate degrader of protein.</p> <p><b>Metabolites produced:</b> Our genomic analysis indicates that most members of this species can produce the following metabolites: acetate, branched chain amino acids, cobalamin (B12), folate (B9), lactate.</p> <p><b>Metabolites consumed:</b> In addition, the genomic analysis indicates that most members of this species do not consume any reported metabolites.</p> <p><b>Emerging Research:</b> This species has been observed at lower levels in individuals with insulin resistance and poor intestinal health. However, despite the likely beneficial role of this species, one study did observe this species at higher levels in patients with autoimmune conditions of the joints.</p>					

# Species Profile

SPECIES

Phylum	Species	Abundance	Range	Level
 Bacteroidota	<i>Alistipes putredinis</i>	0.439%	0.00 - 2.61%	Average
<p>This is a common inhabitant of the gut microbiome.</p> <p><b>Fuel Sources Used:</b> This species is a poor degrader of fibre, a poor degrader of mucin, and a moderate degrader of protein.</p> <p><b>Metabolites produced:</b> Our genomic analysis indicates that most members of this species can produce the following metabolites: butyrate, folate (B9), GABA, lactate, riboflavin (B2), vitamin K.</p> <p><b>Metabolites consumed:</b> In addition, the genomic analysis indicates that most members of this species do not consume any reported metabolites.</p> <p><b>Emerging Research:</b> Studies have observed higher levels of this species in people with some forms of poor intestinal health. However, other studies associated a low abundance of this species with poor intestinal health, so it's role in health is not yet fully understood.</p>				
Proteobacteria	<i>CAG-495 sp000436375</i>	0.432%	0.00 - 0.554%	Average
Firmicutes_A	<i>CAG-269 sp001916065</i>	0.415%	0.00 - 0.00%	High
Firmicutes_A	<i>CAG-103 MIC7540</i>	0.406%	0.00 - 0.418%	Average
Firmicutes_A	<i>CAG-56 sp900066615</i>	0.398%	0.00 - 0.972%	Average
Firmicutes_A	<i>Oscillibacter sp001916835</i>	0.398%	0.00 - 0.377%	High

# Species Profile

## SPECIES

	Phylum	Species	Abundance	Range	Level
+	Firmicutes_A	<i>Coproccoccus_B comes</i>	0.362%	0.086 - 0.813%	Average
<p>This is a common inhabitant of the human gut.</p> <p><b>Fuel Sources Used:</b> This species is a moderate degrader of fibre, a poor degrader of mucin, and a moderate degrader of protein.</p> <p><b>Metabolites produced:</b> Our genomic analysis indicates that most members of this species can produce the following metabolites: acetate, branched chain amino acids, butyrate, cobalamin (B12), folate (B9), lactate, riboflavin (B2), trimethylamine.</p> <p><b>Metabolites consumed:</b> In addition, the genomic analysis indicates that most members of this species do not consume any reported metabolites.</p> <p><b>Emerging Research:</b> This species likely plays a beneficial role in health as it has been observed at lower levels in individuals with poor intestinal health, poor liver health and chronic fatigue syndrome.</p>					
	Firmicutes_A	<i>Ruminococcus_A sp003011855</i>	0.358%	0.00 - 0.892%	Average
	Firmicutes_A	<i>Faecalibacterium prausnitzii_G</i>	0.341%	0.00 - 2.39%	Average
	Firmicutes_A	<i>CAG-83 sp000435975</i>	0.332%	0.00 - 0.382%	Average
	Firmicutes_A	<i>CAG-269 sp003525075</i>	0.316%	0.00 - 0.233%	High
	Firmicutes	<i>CAG-417 sp000432835</i>	0.302%	0.00 - 0.056%	High
	Firmicutes_A	<i>Faecalibacterium prausnitzii_I</i>	0.283%	0.00 - 0.535%	Average

# Species Profile

SPECIES

	Phylum	Species	Abundance	Range	Level
	Firmicutes_A	<i>TF01-11 sp003529475</i>	0.280%	0.00 - 0.396%	Average
	Firmicutes_A	<i>CAG-110 sp000434635</i>	0.259%	0.00 - 0.560%	Average
⊖	Firmicutes	<i>Enterococcus faecalis</i>	0.231%	0.00 - 0.00%	High

This is an inhabitant of the human gut.

**Fuel Sources Used:**

This species is a moderate degrader of fibre, a poor degrader of mucin, and a moderate degrader of protein.

**Metabolites produced:**

Our genomic analysis indicates that most members of this species can produce the following metabolites: acetate, folate (B9), hydrogen sulphide, lactate.

**Metabolites consumed:**

In addition, the genomic analysis indicates that most members of this species do not consume any reported metabolites.

**Emerging Research:**

This species is able to metabolize a medicine called Levodopa, reducing its efficacy. In addition, this species has been associated with complications in people who have undergone intestinal surgery.

# Species Profile

SPECIES

	Phylum	Species	Abundance	Range	Level
+	Actinobacteriota	<i>Bifidobacterium animalis</i>	0.226%	0.00 - 0.365%	Average
<p>This is a naturally occurring human gut bacterium and probiotic; it is sometimes also called <i>Bifidobacterium lactis</i>.</p> <p><b>Fuel Sources Used:</b> This species is a moderate degrader of fibre, mucin, and protein.</p> <p><b>Metabolites produced:</b> Our genomic analysis indicates that most members of this species can produce the following metabolites: acetate, branched chain amino acids, folate (B9), lactate.</p> <p><b>Metabolites consumed:</b> In addition, the genomic analysis shows that most members of this species can consume: oxalate.</p> <p><b>Emerging Research:</b> <i>B. animalis</i> is associated with a low BMI and has been used to improve metabolic disorders such as obesity and poor glucose regulation. It can also produce antimicrobial substances effective against pathogenic bacteria and studies in mouse models suggest it can improve "leaky gut".</p>					
	Firmicutes_A	<i>ER4 sp000765235</i>	0.222%	0.00 - 0.635%	Average

# Species Profile

SPECIES

	Phylum	Species	Abundance	Range	Level
+	Bacteroidota	<i>Odoribacter splanchnicus</i>	0.219%	0.023 - 0.253%	Average
<p>Formerly known as <i>Bacteroides splanchnicus</i>. This a common inhabitant of the human gut.</p> <p><b>Fuel Sources Used:</b> This species is a moderate degrader of fibre, a good degrader of mucin, and a moderate degrader of protein.</p> <p><b>Metabolites produced:</b> Our genomic analysis indicates that most members of this species can produce the following metabolites: acetate, ammonia (urease), biotin (B7), branched chain amino acids, butyrate, folate (B9), GABA, lactate, riboflavin (B2), vitamin K.</p> <p><b>Metabolites consumed:</b> In addition, the genomic analysis shows that most members of this species can consume: GABA.</p> <p><b>Emerging Research:</b> Lower levels of this species have been observed in women with poor glucose regulation and people with poor intestinal health, indicating it likely plays a beneficial role in health.</p>					
	Firmicutes_A	<i>Ruminococcus_D bicirculans</i>	0.218%	0.00 - 3.46%	Average

# Species Profile

## SPECIES

	Phylum	Species	Abundance	Range	Level
⊖	Bacteroidota	<i>Bacteroides_B dorei</i>	0.216%	0.00 - 2.99%	Average
<p>This is a common inhabitant of the gut and is closely related to <i>Bacteroides vulgatus</i>.</p> <p><b>Fuel Sources Used:</b> This species is a good degrader of fibre, a good degrader of mucin, and a moderate degrader of protein.</p> <p><b>Metabolites produced:</b> Our genomic analysis indicates that most members of this species can produce the following metabolites: acetate, beta-glucuronidase, biotin (B7), branched chain amino acids, folate (B9), GABA, lactate, riboflavin (B2), vitamin K.</p> <p><b>Metabolites consumed:</b> In addition, the genomic analysis shows that most members of this species can consume: GABA.</p> <p><b>Emerging Research:</b> Higher levels of this species have been observed in people with poor intestinal health and poor glucose regulation in children. This species has also been associated with diets high in red meat.</p>					
⊖	Euryarchaeota	<i>Methanosphaera stadtmanae</i>	0.209%	0.00 - 0.065%	High
<p>This archaeal species is a methane-producer that can inhabit the human gut.</p> <p><b>Fuel Sources Used:</b> This species is a poor degrader of fibre, a poor degrader of mucin, and a moderate degrader of protein.</p> <p><b>Metabolites produced:</b> Our genomic analysis indicates that most members of this species can produce the following metabolites: acetate, branched chain amino acids, cobalamin (B12), histamine, lactate, methane.</p> <p><b>Metabolites consumed:</b> In addition, the genomic analysis indicates that most members of this species do not consume any reported metabolites.</p> <p><b>Emerging Research:</b> This species can also use the short chain fatty acid acetate as an energy source.</p> <p>Studies have suggested this species can promote inflammation and it has been observed at higher levels in patients with poor intestinal health.</p>					



# Species Profile

## SPECIES

	Phylum	Species	Abundance	Range	Level
	Firmicutes_A	<i>Butyricicoccus_A sp002395695</i>	0.207%	0.00 - 0.202%	High
	Firmicutes_A	<i>Gemmiger MIC8010</i>	0.205%	0.00 - 0.228%	Average
	Firmicutes_A	<i>Agathobacter sp000434275</i>	0.202%	0.00 - 0.043%	High
	Firmicutes_A	<i>Agathobaculum MIC7900</i>	0.201%	0.00 - 0.635%	Average
	Firmicutes_A	<i>Blautia sp000436935</i>	0.199%	0.00 - 0.655%	Average
	Firmicutes_A	<i>Gemmiger MIC9530</i>	0.198%	0.00 - 0.443%	Average
+	Firmicutes_A	<i>Roseburia hominis</i>	0.198%	0.00 - 0.290%	Average

This is a common and important member of the human gut microbiome.

### Fuel Sources Used:

This species is a moderate degrader of fibre, mucin, and protein.

### Metabolites produced:

Our genomic analysis indicates that most members of this species can produce the following metabolites: acetate, beta-glucuronidase, branched chain amino acids, butyrate, folate (B9), hydrogen sulphide, lactate.

### Metabolites consumed:

In addition, the genomic analysis indicates that most members of this species do not consume any reported metabolites.

### Emerging Research:

This is a primary producer of the beneficial short chain fatty acid butyrate. Lower levels of this species have been observed in patients with poor intestinal health and poor heart health. Another study observed higher levels in active women compared to sedentary women. Laboratory studies suggest this species can help regulate the immune system and help reduce inflammation.

# Species Profile

SPECIES

Phylum	Species	Abundance	Range	Level
Firmicutes_A	<i>Eubacterium_F sp000433735</i>	0.195%	0.00 - 0.076%	High
Firmicutes_A	<i>Faecalicatena faecis</i>	0.193%	0.00 - 1.35%	Average
Firmicutes	<i>Merdibacter MIC8457</i>	0.187%	0.00 - 0.070%	High
Firmicutes_A	<i>CAG-83 sp000435555</i>	0.185%	0.00 - 1.33%	Average
Firmicutes_A	<i>F23-B02 sp001916715</i>	0.179%	0.00 - 0.510%	Average
Firmicutes_A	<i>Eubacterium_F sp003491505</i>	0.175%	0.00 - 0.477%	Average
Firmicutes_A	<i>Acetatifactor sp900066565</i>	0.171%	0.00 - 1.29%	Average
Firmicutes_A	<i>UBA11774 sp003507655</i>	0.171%	0.00 - 0.802%	Average
Firmicutes_A	<i>Faecalicatena sp900066545</i>	0.168%	0.00 - 0.046%	High

# Species Profile

SPECIES

	Phylum	Species	Abundance	Range	Level
+	Firmicutes_A	<i>Anaerostipes hadrus</i>	0.167%	0.294 - 5.33%	Low
<p>Formerly known as <i>Eubacterium hadrum</i>. This is a common inhabitant of the human gut.</p> <p><b>Fuel Sources Used:</b> This species is a moderate degrader of fibre, a poor degrader of mucin, and a moderate degrader of protein.</p> <p><b>Metabolites produced:</b> Our genomic analysis indicates that most members of this species can produce the following metabolites: acetate, branched chain amino acids, butyrate, folate (B9), hydrogen sulphide, lactate, riboflavin (B2).</p> <p><b>Metabolites consumed:</b> In addition, the genomic analysis shows that most members of this species can consume: oxalate.</p>					
	Bacteroidota	<i>Alistipes obesi</i>	0.165%	0.00 - 0.430%	Average
	Firmicutes_A	<i>Blautia_A sp000436615</i>	0.160%	0.00 - 1.55%	Average
	Bacteroidota	<i>Barnesiella intestinhominis</i>	0.160%	0.00 - 0.743%	Average
	Proteobacteria	<i>Sutterella wadsworthensis_B</i>	0.159%	0.00 - 0.720%	Average
	Firmicutes_A	<i>Blautia_A MIC9663</i>	0.156%	0.00 - 0.039%	High
	Firmicutes_A	<i>Blautia_A sp900066145</i>	0.145%	0.00 - 0.372%	Average

# Species Profile

## SPECIES

	Phylum	Species	Abundance	Range	Level
+	Bacteroidota	<i>Alistipes shahii</i>	0.143%	0.00 - 0.523%	Average
<p>This is a common inhabitant of the human gut.</p> <p><b>Fuel Sources Used:</b> This species is a moderate degrader of fibre, a good degrader of mucin, and a moderate degrader of protein.</p> <p><b>Metabolites produced:</b> Our genomic analysis indicates that most members of this species can produce the following metabolites: beta-glucuronidase, branched chain amino acids, folate (B9), GABA, lactate, riboflavin (B2).</p> <p><b>Metabolites consumed:</b> In addition, the genomic analysis indicates that most members of this species do not consume any reported metabolites.</p> <p><b>Emerging Research:</b> This species appears to have mostly beneficial effects. It has been associated with beneficial markers of heart health, and was observed at lower levels in people with poor intestinal health. Additionally, a study in mice showed this species may improve the efficacy of some types of immunotherapy. However, this species was also observed at elevated levels in patients with some neurodegenerative conditions.</p>					
+	Firmicutes_A	<i>Faecalibacterium prausnitzii_K</i>	0.139%	0.00 - 2.27%	Average
<p>Previously called <i>Faecalibacterium prausnitzii_A</i> (aka strains SL3/3 and M21/2), this is an inhabitant of the human gut microbiome.</p> <p><b>Fuel Sources Used:</b> This species is a moderate degrader of fibre, a poor degrader of mucin, and a moderate degrader of protein.</p> <p><b>Metabolites produced:</b> Our genomic analysis indicates that most members of this species can produce the following metabolites: acetate, beta-glucuronidase, branched chain amino acids, lactate.</p> <p><b>Metabolites consumed:</b> In addition, the genomic analysis indicates that most members of this species do not consume any reported metabolites.</p> <p><b>Emerging Research:</b> Low levels of <i>F. prausnitzii</i> have been linked to poor intestinal and metabolic health. This species produces lower levels of butyrate than <i>F. prausnitzii_C</i>.</p>					

# Species Profile

SPECIES

	Phylum	Species	Abundance	Range	Level
	Firmicutes	<i>Erysipelatoclostridium sp000752095</i>	0.138%	0.00 - 1.02%	Average
+	Firmicutes_A	<i>Dorea formicigenerans</i>	0.134%	0.107 - 0.418%	Average

Formerly known as *Eubacterium formicigenerans*. This is a common inhabitant of the human gut.

**Fuel Sources Used:**

This species is a poor degrader of fibre, a poor degrader of mucin, and a moderate degrader of protein.

**Metabolites produced:**

Our genomic analysis indicates that most members of this species can produce the following metabolites: acetate, branched chain amino acids, cobalamin (B12), folate (B9), lactate, riboflavin (B2).

**Metabolites consumed:**

In addition, the genomic analysis indicates that most members of this species do not consume any reported metabolites.

**Emerging Research:**

This species has been observed at decreased levels in individuals with poor intestinal health and chronic fatigue syndrome, indicating it likely plays a beneficial role in health.


# Species Profile

SPECIES

	Phylum	Species	Abundance	Range	Level
+	Firmicutes_A	<i>Coprococcus_A catus</i>	0.132%	0.00 - 0.431%	Average
<p>This is an inhabitant of the human gut.</p> <p><b>Fuel Sources Used:</b> This species is a poor degrader of fibre, a poor degrader of mucin, and a moderate degrader of protein.</p> <p><b>Metabolites produced:</b> Our genomic analysis indicates that most members of this species can produce the following metabolites: acetate, branched chain amino acids, cobalamin (B12), folate (B9), lactate, propionate, riboflavin (B2).</p> <p><b>Metabolites consumed:</b> In addition, the genomic analysis indicates that most members of this species do not consume any reported metabolites.</p> <p><b>Emerging Research:</b> Lower levels of this species have been observed in individuals with poor intestinal health and with poor mental health, indicating it likely plays a beneficial role in health.</p>					
	Firmicutes_A	<i>Ruminiclostridium_C sp000435295</i>	0.127%	0.00 - 0.302%	Average
	Firmicutes_A	<i>UBA1417 sp003531055</i>	0.120%	0.00 - 0.982%	Average

# Species Profile

SPECIES

Phylum	Species	Abundance	Range	Level
 Firmicutes_A	<i>Dorea longicatena_B</i>	0.116%	0.00 - 0.747%	Average
<p>This is a common inhabitant of the human gut.</p> <p><b>Fuel Sources Used:</b> This species is a moderate degrader of fibre, a poor degrader of mucin, and a moderate degrader of protein.</p> <p><b>Metabolites produced:</b> Our genomic analysis indicates that most members of this species can produce the following metabolites: acetate, branched chain amino acids, cobalamin (B12), folate (B9), lactate, riboflavin (B2), trimethylamine.</p> <p><b>Metabolites consumed:</b> In addition, the genomic analysis indicates that most members of this species do not consume any reported metabolites.</p> <p><b>Emerging Research:</b> Multiple studies have observed this species at higher levels in individuals with obesity comapred to lean controls. However in obese, postmenopausal women, this species was associated with improved insulin sensitivity. This species has also been observed at lower levels in individuals with chronic fatigue syndrome and poor intestinal health.</p>				
Firmicutes_A	<i>Lachnospiraceae MIC6495</i>	0.113%	0.00 - 0.142%	Average
Firmicutes_A	<i>UBA7160 MIC6745</i>	0.110%	0.00 - 0.308%	Average
Firmicutes_A	<i>Blautia_A sp900120195</i>	0.105%	0.00 - 0.125%	Average
Bacteroidota	<i>Bacteroides thetaiotaomicron</i>	0.102%	0.00 - 0.465%	Average
Firmicutes_A	<i>Faecalibacterium MIC7145</i>	0.102%	0.00 - 1.48%	Average
Firmicutes_A	<i>CAG-273 sp003534295</i>	0.101%	0.00 - 0.272%	Average

# Species Profile

SPECIES

	Phylum	Species	Abundance	Range	Level
	Firmicutes_A	<i>CAG-110 MIC9602</i>	0.097%	0.00 - 0.00%	High
	Firmicutes_A	<i>Ruminococcus_A sp000437095</i>	0.096%	0.00 - 0.253%	Average
+	Firmicutes_A	<i>Eubacterium_I ramulus</i>	0.095%	0.00 - 0.428%	Average

This is an inhabitant of the gut microbiome.

**Fuel Sources Used:**

This species is a moderate degrader of fibre, a poor degrader of mucin, and a moderate degrader of protein.

**Metabolites produced:**

Our genomic analysis indicates that most members of this species can produce the following metabolites: acetate, branched chain amino acids, butyrate, cobalamin (B12), folate (B9), lactate, riboflavin (B2).

**Metabolites consumed:**

In addition, the genomic analysis indicates that most members of this species do not consume any reported metabolites.

**Emerging Research:**

This species can also use the plant flavonoid quercetin as an energy source.

This species has been observed at higher levels in postmenopausal, obese women with insulin resistance. However, another study showed lower levels in patients with poor intestinal health.

	Firmicutes_A	<i>Eubacterium_E hallii_A</i>	0.095%	0.00 - 0.687%	Average
	Actinobacteriota	<i>CAG-1427 sp000436075</i>	0.093%	0.00 - 0.302%	Average
	Firmicutes_A	<i>Faecalibacterium MIC8666</i>	0.092%	0.00 - 0.750%	Average
	Firmicutes_A	<i>Faecalibacterium prausnitzii_A</i>	0.091%	0.00 - 0.455%	Average



# Species Profile

## SPECIES

	Phylum	Species	Abundance	Range	Level
	Firmicutes_A	<i>Lachnospira sp003451515</i>	0.090%	0.00 - 0.400%	Average
	Firmicutes_A	<i>Eisenbergiella sp900066775</i>	0.088%	0.00 - 0.437%	Average
	Firmicutes_A	<i>NK3B98 MIC8311</i>	0.085%	0.00 - 0.048%	High
+	Actinobacteriota	<i>Senegalimassilia anaerobia</i>	0.082%	0.00 - 0.238%	Average

This is a newly discovered inhabitant of the human gut.

**Fuel Sources Used:**

This species is a poor degrader of fibre, a poor degrader of mucin, and a moderate degrader of protein.

**Metabolites produced:**

Our genomic analysis indicates that most members of this species can produce the following metabolites: acetate, branched chain amino acids, lactate, riboflavin (B2), vitamin K.

**Metabolites consumed:**

In addition, the genomic analysis indicates that most members of this species do not consume any reported metabolites.

**Emerging Research:**

Lower levels of this species have been observed in individuals with poor intestinal health, suggesting it may play a beneficial role in health.

	Firmicutes_A	<i>CAG-170 MIC9129</i>	0.081%	0.00 - 0.077%	High
	Firmicutes_A	<i>UBA7597 MIC7484</i>	0.080%	0.00 - 0.060%	High
	Firmicutes_A	<i>Lachnospira sp000436475</i>	0.074%	0.00 - 0.141%	Average

# Species Profile

SPECIES

Phylum	Species	Abundance	Range	Level
Firmicutes_A	<i>Agathobaculum butyriciproducens</i>	0.073%	0.00 - 0.621%	Average
Firmicutes_A	<i>GCA-900066135 MIC6659</i>	0.073%	0.00 - 0.247%	Average
Firmicutes_A	<i>ER4 sp900317525</i>	0.073%	0.00 - 0.375%	Average
Firmicutes_A	<i>CAG-74 MIC7044</i>	0.070%	0.00 - 0.260%	Average
Firmicutes_A	<i>CAG-110 sp003525905</i>	0.070%	0.00 - 0.331%	Average
Actinobacteriota	<i>Enorma massiliensis</i>	0.070%	0.00 - 0.00%	High
Firmicutes_A	<i>Hungatella_A MIC8772</i>	0.068%	0.00 - 0.077%	Average
Firmicutes_A	<i>CAG-81 sp900066535</i>	0.067%	0.00 - 0.164%	Average
Firmicutes_A	<i>Oscillibacter MIC7608</i>	0.067%	0.00 - 0.067%	Average
Firmicutes_A	<i>Clostridium_Q sp003024715</i>	0.066%	0.00 - 0.265%	Average
Firmicutes_A	<i>Ruminococcaceae MIC8156</i>	0.066%	0.00 - 0.058%	High
Firmicutes_A	<i>Angelakisella MIC6791</i>	0.064%	0.00 - 0.305%	Average
Bacteroidota	<i>Bacteroides_A MIC8590</i>	0.064%	0.00 - 0.00%	High

# Species Profile

SPECIES

	Phylum	Species	Abundance	Range	Level
	Firmicutes_A	<i>Eubacterium_F MIC7117</i>	0.063%	0.00 - 0.00%	High
	Firmicutes_A	<i>CAG-170 sp003516765</i>	0.059%	0.00 - 0.087%	Average
⊖	Bacteroidota	<i>Parabacteroides merdae</i>	0.057%	0.00 - 0.482%	Average

Formerly known as *Bacteroides merdae*. This is a common inhabitant of the human gut.

**Fuel Sources Used:**

This species is a moderate degrader of fibre, a good degrader of mucin, and a moderate degrader of protein.

**Metabolites produced:**

Our genomic analysis indicates that most members of this species can produce the following metabolites: acetate, beta-glucuronidase, biotin (B7), branched chain amino acids, folate (B9), GABA, lactate, riboflavin (B2), vitamin K.

**Metabolites consumed:**

In addition, the genomic analysis shows that most members of this species can consume: GABA.

**Emerging Research:**

Higher levels of this species have been observed in individuals with elevated blood pressure, and poor intestinal health. This species has been associated with a diet low in fruits and vegetables.

	Actinobacteriota	<i>Adlercreutzia MIC8014</i>	0.057%	0.00 - 0.269%	Average
	Firmicutes_A	<i>UBA7160 MIC9207</i>	0.057%	0.00 - 0.188%	Average
	Firmicutes_A	<i>QALW01 MIC8348</i>	0.057%	0.00 - 0.00%	High
	Firmicutes_A	<i>Eubacterium_F MIC7373</i>	0.054%	0.00 - 0.00%	High

# Species Profile

SPECIES

Phylum	Species	Abundance	Range	Level
Firmicutes_A	<i>Eubacterium_E MIC8705</i>	0.051%	0.00 - 0.00%	High
Proteobacteria	<i>Parasutterella excrementihominis</i>	0.051%	0.00 - 0.389%	Average
Firmicutes_A	<i>UBA7182 MIC8422</i>	0.050%	0.00 - 0.090%	Average
Firmicutes_A	<i>CAG-81 sp900066785</i>	0.048%	0.00 - 0.146%	Average
Firmicutes_A	<i>CAG-272 MIC7215</i>	0.046%	0.00 - 0.075%	Average
Firmicutes_A	<i>GCA-900066575 MIC7948</i>	0.046%	0.00 - 0.093%	Average
Firmicutes_A	<i>Blautia_A sp900066505</i>	0.045%	0.00 - 0.092%	Average

# Species Profile

SPECIES

	Phylum	Species	Abundance	Range	Level
⊖	Firmicutes	<i>Streptococcus salivarius</i>	0.045%	0.00 - 0.306%	Average
<p>This is a common inhabitant of the human oral microbiota and is also found in the human gut.</p> <p><b>Fuel Sources Used:</b> This species is a moderate degrader of fibre, a poor degrader of mucin, and a moderate degrader of protein.</p> <p><b>Metabolites produced:</b> Our genomic analysis indicates that most members of this species can produce the following metabolites: acetate, ammonia (urease), branched chain amino acids, folate (B9), lactate.</p> <p><b>Metabolites consumed:</b> In addition, the genomic analysis indicates that most members of this species do not consume any reported metabolites.</p> <p><b>Emerging Research:</b> This species has been observed at higher levels in patients with poor heart health and poor intestinal health, however one study observed it at lower levels in people with a different intestinal health condition.</p>					
	Firmicutes_A	<i>Blautia_A MIC8343</i>	0.041%	0.00 - 0.287%	Average
	Firmicutes_A	<i>Eubacterium_G MIC6759</i>	0.040%	0.00 - 0.00%	High
	Firmicutes_A	<i>UBA9502 MIC6887</i>	0.040%	0.00 - 0.080%	Average
	Firmicutes_A	<i>Agathobacter sp900317585</i>	0.038%	0.00 - 0.00%	High
	Firmicutes_A	<i>CAG-81 sp000435795</i>	0.036%	0.00 - 0.104%	Average
	Eukaryote_unclassified	<i>Blastocystis sp. subtype 4</i>	0.035%	0.00 - 0.00%	High

# Species Profile

SPECIES

Phylum	Species	Abundance	Range	Level
Firmicutes_A	<i>Blautia_A MIC7077</i>	0.035%	0.00 - 0.126%	Average
Firmicutes_A	<i>GCA-900066135 sp900066135</i>	0.033%	0.00 - 0.086%	Average
Firmicutes_A	<i>Blautia_A MIC7810</i>	0.033%	0.00 - 0.083%	Average
Firmicutes_A	<i>CAG-81 sp900066055</i>	0.033%	0.00 - 0.035%	Average
Firmicutes_A	<i>Eubacterium_I ramulus_A</i>	0.033%	0.00 - 0.140%	Average
Firmicutes_A	<i>Blautia_A MIC8910</i>	0.032%	0.00 - 0.059%	Average
Firmicutes_A	<i>Marvinbryantia sp900066075</i>	0.031%	0.00 - 0.062%	Average
Firmicutes_A	<i>Eubacterium_G sp000432355</i>	0.031%	0.00 - 0.057%	Average
Firmicutes_A	<i>Clostridium_M sp000431375</i>	0.030%	0.00 - 0.218%	Average
Firmicutes_A	<i>Blautia_A sp000285855</i>	0.030%	0.00 - 0.129%	Average
Firmicutes_A	<i>Dorea sp900066765</i>	0.028%	0.00 - 0.032%	Average

# Species Profile

## SPECIES

	Phylum	Species	Abundance	Range	Level
⊖	Bacteroidota	<i>Alistipes_A indistinctus</i>	0.025%	0.00 - 0.064%	Average
<p>This is a recently discovered inhabitant of the human gut.</p> <p><b>Fuel Sources Used:</b> This species is a moderate degrader of fibre, mucin, and protein.</p> <p><b>Metabolites produced:</b> Our genomic analysis indicates that most members of this species can produce the following metabolites: acetate, branched chain amino acids, GABA, lactate, riboflavin (B2).</p> <p><b>Metabolites consumed:</b> In addition, the genomic analysis indicates that most members of this species do not consume any reported metabolites.</p> <p><b>Emerging Research:</b> High levels of this species have been observed in individuals with poor glucose regulation. However one study observed reduced levels in individuals with poor intestinal health.</p>					
	Firmicutes_A	<i>Lachnospiraceae MIC8879</i>	0.025%	0.00 - 0.128%	Average
	Firmicutes_A	<i>CAG-272 sp000433515</i>	0.024%	0.00 - 0.033%	Average
	Firmicutes_A	<i>Dorea sp000433215</i>	0.024%	0.00 - 0.077%	Average
	Firmicutes_A	<i>CAG-272 MIC9439</i>	0.023%	0.00 - 0.00%	High
	Firmicutes_A	<i>UBA1390 MIC8736</i>	0.022%	0.00 - 0.00%	High

# Species Profile

SPECIES

	Phylum	Species	Abundance	Range	Level
⊖	Desulfobacterota_	<i>Bilophila wadsworthia</i>	0.022%	0.00 - 0.200%	Average
	A				
<p>This is a common inhabitant of the human gut, but can become problematic at high levels.</p> <p><b>Fuel Sources Used:</b> This species is a poor degrader of fibre, a poor degrader of mucin, and a moderate degrader of protein.</p> <p><b>Metabolites produced:</b> Our genomic analysis indicates that most members of this species can produce the following metabolites: acetate, ammonia (urease), branched chain amino acids, hydrogen sulphide, lactate, riboflavin (B2), trimethylamine.</p> <p><b>Metabolites consumed:</b> In addition, the genomic analysis indicates that most members of this species do not consume any reported metabolites.</p> <p><b>Emerging Research:</b> Higher levels of this species have been observed in people with poor intestinal health and in people that have a diet high in saturated fats. Mice studies have also suggested this species can promote increased inflammation in the gut and increased barrier dysfunction, though more research needs to be conducted in humans to confirm these results.</p>					
	Firmicutes_A	<i>UBA7182 MIC8257</i>	0.022%	0.00 - 0.036%	Average
	Firmicutes_A	<i>Lawsonibacter asaccharolyticus</i>	0.022%	0.00 - 0.127%	Average
	Firmicutes_A	<i>Monoglobaceae MIC9391</i>	0.020%	0.00 - 0.00%	High
	Desulfobacterota_	<i>Mailhella sp003150275</i>	0.020%	0.00 - 0.019%	High
	A				
	Firmicutes_A	<i>Marvinbryantia MIC9792</i>	0.020%	0.00 - 0.077%	Average
	Firmicutes_A	<i>Oscillospiraceae MIC8045</i>	0.019%	0.00 - 0.042%	Average



# Species Profile

SPECIES

Phylum	Species	Abundance	Range	Level
Firmicutes_A	<i>QALS01 MIC9566</i>	0.019%	0.00 - 0.035%	Average
Firmicutes	<i>CAG-313 sp000433035</i>	0.019%	0.00 - 0.229%	Average
Firmicutes_A	<i>CAG-269 sp001916005</i>	0.019%	0.00 - 0.053%	Average
Bacteroidota	<i>Alistipes_A sp900240235</i>	0.018%	0.00 - 0.069%	Average
Firmicutes_A	<i>UBA9502 sp003506385</i>	0.018%	0.00 - 0.00%	High
Firmicutes_A	<i>Butyricicoccaceae MIC8222</i>	0.017%	0.00 - 0.023%	Average
Firmicutes_A	<i>CAG-110 MIC9052</i>	0.017%	0.00 - 0.094%	Average
Bacteroidota	<i>Coprobacter secundus</i>	0.016%	0.00 - 0.031%	Average
Firmicutes_A	<i>Anaerovoracaceae MIC7161</i>	0.014%	0.00 - 0.011%	High
Firmicutes_A	<i>Lachnospiraceae MIC6593</i>	0.014%	0.00 - 0.036%	Average
Firmicutes_A	<i>Oscillibacter MIC7446</i>	0.013%	0.00 - 0.033%	Average
Firmicutes_A	<i>CAG-269 sp000431335</i>	0.012%	0.00 - 0.096%	Average
Firmicutes_A	<i>Faecalicatena MIC9341</i>	0.011%	0.00 - 0.00%	High

# Species Profile

SPECIES

	Phylum	Species	Abundance	Range	Level
+	Firmicutes	<i>Lactobacillus_C rhamnosus</i>	0.010%	0.00 - 0.00%	High
<p>This species is commonly found in fermented dairy products but can also be found in the human gut.</p> <p><b>Fuel Sources Used:</b> This species is a moderate degrader of fibre, a poor degrader of mucin, and a moderate degrader of protein.</p> <p><b>Metabolites produced:</b> Our genomic analysis indicates that most members of this species can produce the following metabolites: acetate, beta-glucuronidase, folate (B9), lactate.</p> <p><b>Metabolites consumed:</b> In addition, the genomic analysis indicates that most members of this species do not consume any reported metabolites.</p> <p><b>Emerging Research:</b> Studies in children have shown some strains may help reduce diarrhoea, respiratory infections, and abdominal pain associated with poor intestinal health. In general, Lactobacillus species do not colonise the adult human gut and are only transient.</p>					



# Gut microbiome report

