
Dietary considerations for increasing fibre intake

Adequate intake (AI) of dietary fibre in children




Suggested dietary target (SDT) for fibre intake to reduce risk of chronic disease.¹


Women 28g


Men 38g


Al and SDT fibre intake for adults





Example of a meal plan to meet daily fibre requirements











BRISTOL STOOL CHART ²					
Type 1		Separate hard lumps like nuts (difficult to pass)			
Type 2		Sausage-shaped but lumpy			
Type 3		Like a sausage, but with cracks on the surface			
Type 4		Like a sausage or snake, smooth and soft			
Type 5		Soft blobs with clear-cut edges (passed easily)			
Type 6		Fluffy pieces with ragged edges, a mushy stool			
Type 7		Watery, no solid pieces (entirely liquid)			

The following table gives approximate values for soluble, insoluble and total fibre content of foods commonly eaten as part of a varied, healthy, predominantly wholefood diet.

Also included is the % daily value (%DV); based on suggested dietary target (SDT) for fibre intake: **women= 28g, men= 38g**.

FOOD	SERVING SIZE	SOLUBLE FIBRE (G)	INSOLUBLE FIBRE (G)	TOTAL FIBRE (G)	%DV
Lentils, cooked	½ cup	3.3	4.5	7.8	Women: 28%
Pinto beans, cooked	½ cup	5.5	1.9	7.4	Men: 21% Women: 26%
Artichoke (globe or	½ cup	5.2	2.0	7.2	Men: 19% Women: 26%
french), hearts cooked Black beans, cooked	½ cup	3.8	3.1	6.9	Men 19% Women: 25%
Wholegrain pasta, cooked	1 cup	4.1	2.2	6.3	Men: 18% Women: 23%
Kidney beans, cooked	½ cup	2.9	2.9	5.8	Men: 17% Women: 21%
Pear, with skin	1 medium	1.1	4.4	5.5	Men: 15% Women: 20%
Broccoli, cooked	1 cup	2.6	2.9	5.5	Men: 14% Women: 20%
Apple, with skin	1 medium	3.2	1.2	4.4	Men: 14% Women: 16%
					Men: 12% Women: 16%
Green peas, cooked	½ cup	3.2	1.2	4.4	Men: 12% Women: 15%
Lima beans, cooked	½ cup	2.1	2.2	4.3	Men 11% Women: 15%
Barley, cooked	½ cup	3.3	0.9	4.2	Men: 11% Women: 14%
Okra, cooked	½ cup	3.1	1.0	4.0	Men: 11% Women: 14%
Oatmeal, cooked	1 cup	2.4	1.6	4.0	Men: 11% Women: 14%
Passion fruit, purple, pulp	2 fruits	2.0	1.8	3.8	Women: 14% Men: 10% Women: 14%
Blackberries	½ cup	3.1	0.07	3.8	Men: 10%
Brussel sprouts, cooked	1 cup	1.7	1.9	3.6	Women: 13% Men: 9%
Orange	1 medium	2.1	1.3	3.4	Women: 12% Men: 9%
Kiwi fruit	1 large	2.4	0.8	3.2	Women: 11% Men: 8%
Raspberries	½ cup	0.9	2.3	3.2	Women: 11% Man: 8%
Prunes, dried	4 medium	1.3	1.8	3.1	Women: 11% Men: 8%
Strawberries	1 cup	1.2	1.8	3.0	Women: 11% Men: 8%
Sunflower seeds, kernels, dried	1/4 cup	1.1	1.9	3.0	Women: 11% Men: 8%
Potato with skin, baked	1 medium	1.5	1.5	3.0	Women: 11% Men: 8%
Wholegrain bread	1 slice	2.8	0.1	2.9	Women: 10% Men: 8%
Banana	1 medium	2.1	0.7	2.8	Women: 10% Men: 7%
Quinoa, cooked	½ cup	1.1	1.5	2.6	Women: 9% Men: 7%
Carrots, raw	1 medium	1.1	1.5	2.6	Women: 9% Men: 7%
Kale, chopped, cooked	1 cup	0.8	1.8	2.6	Women: 9% Men: 7%
Zucchini, sliced, cooked	1 cup	1.3	1.2	2.5	Women: 9% Men: 7%
Flaxseeds	1 tbsp.	1.3	1.1	2.4	Women: 9% Men: 6%
Figs, dried	3 (24g)	1.4	1.0	2.4	Women: 9%
Avocado	1/4 (50g)	1.3	0.8	2.1	Men: 6% Women: 8%
Blueberries	½ cup	0.85	1.25	2.1	Men: 6% Women: 8%
Sweet potato with skin,	1 small	1.1	0.9	2.0	Men: 6% Women: 7%
baked Green string beans,	½ cup	0.8	1.2	2.0	Men: 5% Women: 7%
cooked	2 fruits	1.0	0.8	1.8	Men: 5% Women: 6%
			1.2		Men: 5% Women: 5%
Almonds, raw	10 kernels	0.2		1.4	Men: 4% Women: 5%
Brown rice, cooked	½ cup	1.3	0.1	1.4	Men: 4% Women: 5%
Walnuts, raw, shelled	10 halves	0.2	1.1	1.3	Men: 3% Women: 5%
Tomato with skin	1 medium	0.3	1.0	1.3	Men: 3%

References: 1. https://www.nrv.gov.au/chronic-disease/summary 2. World Gastroenterology Organisation (WGO) Global Guidelines. (2010). Constipation: a global perspective

